

# THE $Q$ - $q$ -SCHUR SUPERALGEBRA

JIE DU AND JINKUI WAN<sup>0</sup>

**ABSTRACT.** As a natural generalisation of  $q$ -Schur algebras associated with the Hecke algebra  $\mathcal{H}_{r,R}$  (of the symmetric group), we introduce the  $Q$ - $q$ -Schur superalgebra associated with the Hecke-Clifford superalgebra  $\mathcal{H}_{r,R}^c$ , which, by definition, is the endomorphism algebra of the induced  $\mathcal{H}_{r,R}^c$ -module from certain  $q$ -permutation modules over  $\mathcal{H}_{r,R}$ . We will describe certain integral bases for these superalgebras in terms of matrices and will establish the base change property for them. We will also identify the  $Q$ - $q$ -Schur superalgebras with the quantum queer Schur superalgebras investigated in the context of quantum queer supergroups and then provide a classification of their irreducible representations over a certain extension of the field  $\mathbb{C}(\mathbf{v})$  of complex rational functions.

*We dedicate the paper to the memory of Professor James A. Green*

## 1. INTRODUCTION

In the determination of the polynomial representations of the complex general linear group  $GL_n(\mathbb{C})$ , Issai Schur introduced certain finite dimensional algebras to describe the homogeneous components of these representations and to set up a correspondence between the representations of  $GL_n(\mathbb{C})$  of a fixed homogeneous degree  $r$  and the representations of the symmetric group  $\mathfrak{S}_r$  on  $r$  letters. This correspondence forms part of the well-known Schur–Weyl duality. J.A. Green publicised this algebraic approach in his 1980 monograph [14] and presented the theory “in some ways very much in keeping with the present-day ideas on representations of algebraic groups”. In particular, he used the term *Schur algebras* for these finite dimensional algebras and provided some standard combinatorial treatment for the structure and representations of Schur algebras.

Since the introduction of quantum groups in mid 1980s, the family of Schur algebras has been enlarged to include  $q$ -Schur algebras, affine  $q$ -Schur algebras, and their various super analogues. See, e.g., [7], [5], [4], [11] and the references therein. Thus, the treatment in [14] has also been “quantised” in various forms which provide applications to both structure and representations of the associated quantum linear groups, finite general linear groups, general linear  $p$ -adic groups, general linear supergroups and so on. In this paper, we will further generalise Green’s standard treatment to a new class of Schur superalgebras, the  $Q$ - $q$ -Schur superalgebra.

Originally arising from the classification of complex simple superalgebras, there are two types of  $q$ -Schur superalgebras. The  $q$ -Schur superalgebra of type  $M$  links representations of quantum linear supergroups with those of Hecke algebras of type  $A$ , while the  $q$ -Schur superalgebra of type  $Q$  links the quantum queer supergroup with the Hecke–Clifford superalgebra. It is known that the former has various structures shared with the  $q$ -Schur

---

<sup>0</sup>Corresponding author.

*Date:* November 18, 2015.

Supported by ARC DP-120101436 and NSFC-11101031. The research was carried out while Wan was visiting the University of New South Wales during the year 2012–2013. The hospitality and support of UNSW are gratefully acknowledged.

algebra; see [19], [11], [13] and [9], while the latter has been introduced in [20, 12] in the context of Schur–Weyl–Sergeev duality and investigated in [12] in terms of a Drinfeld–Jimbo type presentation (cf. [13]).

In this paper, we will explore the combinatorial properties of the  $q$ -Schur superalgebra of type  $\mathbf{Q}$ . Thus, like the  $q$ -Schur algebra, we first introduce a new class of superalgebras, called the  $\mathbf{Q}$ - $q$ -Schur superalgebra, which is defined as the endomorphism algebra of a direct sum of certain “ $q$ -permutation supermodules” for the Hecke–Clifford superalgebra. We then construct its standard basis arising from analysing double cosets in the symmetric group. In particular, we use matrices to label the basis elements. In this way, we establish the base change property for the  $\mathbf{Q}$ - $q$ -Schur superalgebra. We then identify them in the generic case with the quotients of the quantum queer supergroups and classify their irreducible representations in this case.

Here is the layout of the paper. In §2, we review the definition of the Hecke–Clifford superalgebra  $\mathcal{H}_{r,R}^c$  over a commutative ring  $R$  and its standard basis. We then introduce certain (induced)  $q$ -permutation supermodules and define the  $\mathbf{Q}$ - $q$ -Schur superalgebra  $\mathcal{Q}_q(n, r; R)$ . In §3, we study the properties of  $q$ -permutations supermodules and investigate some new bases. In §4, we introduce some special elements in the Clifford superalgebras which is the key to a basis for the endomorphism algebra of the induced  $q$ -trivial representation. An integral basis and the base change property for the  $\mathbf{Q}$ - $q$ -Schur superalgebra are given in §5. We then identify the  $\mathbf{Q}$ - $q$ -Schur superalgebras as quotients of the quantum queer supergroup in §6. Using the identification, we establish a classification of irreducible representations of the  $\mathbf{Q}$ - $q$ -Schur superalgebra in the last section.

*Throughout the paper*, let  $\mathbb{Z}_2 = \{0, 1\}$ . We will use a twofold meaning for  $\mathbb{Z}_2$ . We will regard  $\mathbb{Z}_2$  as an abelian group when we use it to describe a superspace. However, for a matrix or an  $n$ -tuple with entries in  $\mathbb{Z}_2$ , we will regard it as a subset of  $\mathbb{Z}$ .

Let  $\mathcal{A} = \mathbb{Z}[\mathbf{q}]$  and let  $R$  be a commutative ring which is also an  $\mathcal{A}$ -module by mapping  $\mathbf{q}$  to  $q \in R$ . We assume that the characteristic of  $R$  is not equal to 2. If  $\mathcal{Q}$  denotes an  $\mathcal{A}$ -algebra or an  $\mathcal{A}$ -module, then  $\mathcal{Q}_R = \mathcal{Q} \otimes_{\mathcal{A}} R$  denotes the object obtained by base change to  $R$ . From §6 on wards, we will use the ring  $\mathcal{Z} = \mathbb{Z}[\mathbf{v}, \mathbf{v}^{-1}]$  of Laurent polynomials in indeterminate  $\mathbf{v}$ , where  $\mathbf{q} = \mathbf{v}^2$ .

## 2. HECKE-CLIFFORD SUPERALGEBRAS AND $\mathbf{Q}$ - $q$ -SCHUR SUPERALGEBRAS

We first introduce an endomorphism algebra of an induced module from certain “ $q$ -permutation modules” over the Hecke–Clifford superalgebra. We will prove in §6 that it is isomorphic to the quantum queer Schur superalgebra defined in [12] by the tensor space of the natural representation of the quantum queer supergroup  $U_{\mathbf{v}}(\mathfrak{q}_n)$ .

Let  $R$  be a commutative ring of characteristic not equal to 2 and let  $q \in R$ . Let  $\mathcal{C}_r$  denote the *Clifford superalgebra* over  $R$  generated by odd elements  $c_1, \dots, c_r$  subject to the relations

$$c_i^2 = -1, \quad c_i c_j = -c_j c_i, \quad 1 \leq i \neq j \leq r. \quad (2.0.1)$$

The *Hecke-Clifford superalgebra*  $\mathcal{H}_{r,R}^c$  is the associative superalgebra over the ring  $R$  with the even generators  $T_1, \dots, T_{r-1}$  and the odd generators  $c_1, \dots, c_r$  subject to (2.0.1) and the following additional relations: for  $1 \leq i, i' \leq r-1$  and  $1 \leq j \leq r$  with  $|i - i'| > 1$  and  $j \neq i, i+1$ ,

$$\begin{aligned} (T_i - q)(T_i + 1) &= 0, & T_i T_{i'} &= T_{i'} T_i, & T_i T_{i+1} T_i &= T_{i+1} T_i T_{i+1}, \\ T_i c_j &= c_j T_i, & T_i c_i &= c_{i+1} T_i, & T_i c_{i+1} &= c_i T_i - (q - 1)(c_i - c_{i+1}). \end{aligned} \quad (2.0.2)$$

Note that if  $q$  is invertible then  $T_i^{-1}$  exists and the last relation is obtained by taking the inverses of the last second relation.

For notational simplicity, if  $R = \mathcal{A} := \mathbb{Z}[\mathbf{q}]$ , the integral polynomial ring in  $\mathbf{q}$ , then we write  $\mathcal{H}_r^c = \mathcal{H}_{r,\mathcal{A}}^c$ .

Let  $\mathfrak{S}_r$  be the symmetric group on  $\{1, 2, \dots, r\}$  which is generated by the simple transpositions  $s_k = (k, k+1)$  for  $1 \leq k \leq r-1$ . The subalgebra  $\mathcal{H}_{r,R}$  of  $\mathcal{H}_{r,R}^c$  generated by  $T_1, \dots, T_{r-1}$  is the (Iwahori-)Hecke algebra associated with  $\mathfrak{S}_r$ .

**Remark 2.1.** Jones and Nazarov [16] introduced the notion of an affine Hecke-Clifford superalgebra which is generated by  $T_1, \dots, T_{r-1}, c_1, \dots, c_r, X_1, \dots, X_r$  subject to the relations (2.0.1), (2.0.2) and certain additional explicit relations which are not mentioned here. It is known from [16, Proposition 3.5] (cf. also [1, Remark 4.2]) that  $\mathcal{H}_{r,R}^c$  can naturally be viewed as a quotient superalgebra of the affine Hecke-Clifford superalgebra by the two-sided ideal generated by  $X_1$ .

We also note that the generator  $T_i$  used in [1] is equal to  $v^{-1}T_i$  here, where  $v^2 = q$  for some  $v \in R$ . Moreover, we choose  $c_j^2 = -1$  in (2.0.1), following [20, 21], whereas in *loc. cit.* the authors take  $C_j^2 = 1$ . Thus, assuming  $\sqrt{-1} \in R$ , we have  $c_i = \sqrt{-1}C_i$ .

Like the Hecke algebra  $\mathcal{H}_{r,R}$ , the symmetry of  $\mathcal{H}_{r,R}^c$  can be seen from the following (anti-)automorphisms of order 2.

**Lemma 2.2.** *The  $R$ -algebra  $\mathcal{H}_{r,R}^c$  admits the following algebra involutions  $\varphi, \psi$  and anti-involutions  $\tau, \iota$  defined by*

- (1)  $\varphi : T_i \mapsto -T_{r-i} + (q-1), \quad c_j \mapsto c_{r+1-j};$
- (2)  $\psi : T_i \mapsto c_i T_i c_{i+1}, \quad c_j \mapsto c_j;$
- (3)  $\tau : T_i \mapsto -T_i + (q-1), \quad c_j \mapsto c_j;$
- (4)  $\iota : T_i \mapsto T_i - (q-1)c_i c_{i+1}, \quad c_j \mapsto c_j,$

for all  $1 \leq i, j \leq r$  and  $i \neq r$ ,

*Proof.* Note that the correspondence  $T_i \mapsto -T_i + (q-1)$  in (3), denoted by  $(\ )^\#$  in [7], defines an automorphism of the subalgebra  $\mathcal{H}_{r,R}$ . So (3) follows easily. (If  $q^{-1} \in R$ , then (3) is seen easily by the fact  $-T_i + (q-1) = -qT_i^{-1}$ .) By directly checking the relations (2.0.1) and (2.0.2), there is an anti-involution

$$\gamma : T_i \longmapsto T_{r-i}, c_j \longmapsto c_{r+1-j}.$$

Thus, compositing  $\gamma$  and  $\tau$  gives (1).

Let  $\dot{T}_i = c_i T_i c_{i+1}$ . Then  $\dot{T}_i, c_j$  generate  $\mathcal{H}_{r,R}^c$  and satisfy all the relations (2.0.1) and (2.0.2) above with  $T_i$  replaced by  $\dot{T}_i$ . Hence,  $\psi$  in (2) is an automorphism of order 2. Finally, the map  $\iota$  in (4) is the map  $\psi$  followed by the map  $\tau$  defined relative to the generators  $\dot{T}_i, c_j$ , since, by the last relation in (2.0.2),

$$\iota(T_i) = T_i - (q-1)c_i c_{i+1} = -c_i T_i c_{i+1} + (q-1).$$

(Of course, by noting Remark 2.1, assertion (4) follows also from [1, (2.31)].)  $\square$

Let  $T_{s_i} = T_i$  and  $T_w = T_{s_{i_1}} \cdots T_{s_{i_k}}$  where  $w = s_{i_1} \cdots s_{i_k} \in \mathfrak{S}_r$  is a reduced expression. Let  $\leq$  be the Bruhat order of  $\mathfrak{S}_r$ , that is,  $\pi \leq w$  if  $\pi$  is a subexpression of some reduced expression for  $w$ . For  $\alpha = (\alpha_1, \dots, \alpha_r) \in \mathbb{Z}_2^r$ , set  $c^\alpha = c_1^{\alpha_1} \cdots c_r^{\alpha_r}$ . Observe that the symmetric group  $\mathfrak{S}_r$  acts naturally on  $\mathcal{C}_r$  by permutating the generators  $c_1, \dots, c_r$ . We shall denote this action by  $z \mapsto w \cdot z$  for  $z \in \mathcal{C}_r, w \in \mathfrak{S}_r$ . Then we have

$$w \cdot c^\alpha = c_{w(1)}^{\alpha_1} \cdots c_{w(r)}^{\alpha_r} = \pm c^{\alpha \cdot w^{-1}},$$

where  $\alpha \cdot w = (\alpha_{w(1)}, \dots, \alpha_{w(r)})$ , for  $\alpha \in \mathbb{Z}_2^r$ ,  $w \in \mathfrak{S}_r$ , is the right place permutation action. Moreover, by (2.0.2), we have

$$c_k^{\alpha_k} c_{k+1}^{\alpha_{k+1}} T_{s_k} = T_{s_k} c_{k+1}^{\alpha_k} c_k^{\alpha_{k+1}} + (q-1)(c_k^{\alpha_k + \alpha_{k+1}} - c_{k+1}^{\alpha_k} c_k^{\alpha_{k+1}}) \quad (2.2.1)$$

Thus, by induction on the length  $\ell(w)$  of  $w$ , one proves the following formula: for any  $\alpha \in \mathbb{Z}_2^r$  and  $w \in \mathfrak{S}_r$

$$c^\alpha T_w = T_w(w^{-1} \cdot c^\alpha) + \sum_{\pi < w, \beta \in \mathbb{Z}_2^r} a_{\pi, \beta}^{\alpha, w} T_\pi c^\beta = \pm T_w c^{\alpha \cdot w} + \sum_{\pi < w, \beta \in \mathbb{Z}_2^r} a_{\pi, \beta}^{\alpha, w} T_\pi c^\beta, \quad (2.2.2)$$

for some  $a_{\pi, \beta}^{\alpha, w} \in R$ . Then, by the results [16, Proposition 2.1] and [1, Theorem 3.6] over fields, it is easy to deduce that the following holds.

**Lemma 2.3.** *The superalgebra  $\mathcal{H}_{r,R}^c$  is a free  $R$ -module. Moreover, both the sets  $\mathcal{B} = \{T_w c^\alpha \mid w \in \mathfrak{S}_r, \alpha \in \mathbb{Z}_2^r\}$  and  $\{c^\alpha T_w \mid w \in \mathfrak{S}_r, \alpha \in \mathbb{Z}_2^r\}$  form bases for  $\mathcal{H}_{r,R}^c$ .*

*Proof.* By (2.2.2), it suffices to prove that  $\mathcal{B}$  forms a basis. First, the set spans  $\mathcal{H}_{r,R}^c$  by (2.2.2). Suppose next  $R = \mathcal{A}$ . By base change to the fraction field of  $\mathcal{A}$  and [1, Theorem 3.6], we see that  $\mathcal{B}$  is linearly independent and, hence, forms an  $\mathcal{A}$ -basis for  $\mathcal{H}_r^c$ . For a general  $R$ , specialization to  $R$  by sending  $\mathbf{q}$  to  $q$  gives a basis  $\{T_w c^\alpha \otimes 1 \mid w \in \mathfrak{S}_r, \alpha \in \mathbb{Z}_2^r\}$  for  $\mathcal{H}_r^c \otimes_{\mathcal{A}} R$ . Now, the presentation of  $\mathcal{H}_{r,R}^c$  gives an  $R$ -algebra epimorphism  $f : \mathcal{H}_{r,R}^c \rightarrow \mathcal{H}_r^c \otimes_{\mathcal{A}} R$ , sending  $T_i, c_j$  to  $T_i \otimes 1, c_j \otimes 1$ , respectively, and sending  $\mathcal{B}$  to a basis. This forces that  $\mathcal{B}$  itself must be a basis (and that  $f$  is an isomorphism).  $\square$

**Corollary 2.4.** *Let  $R$  be a commutative ring which is an  $\mathcal{A}$ -module via the map  $\mathcal{A} \rightarrow R, \mathbf{q} \mapsto q$ .*

- (1) *There is an algebra isomorphism  $\mathcal{H}_{r,R}^c \cong \mathcal{H}_r^c \otimes_{\mathcal{A}} R$ .*
- (2) *As a module over the subalgebra  $\mathcal{H}_{r,R}$ ,  $\mathcal{H}_{r,R}^c$  is  $\mathcal{H}_{r,R}$ -free with basis  $\{c^\alpha \mid \alpha \in \mathbb{Z}_2^r\}$ .*

*Proof.* The proof for (1) is given above. Observe that the  $R$ -submodule  $\mathcal{H}_{r,R} c^\alpha$  of  $\mathcal{H}_{r,R}^c$  is spanned by the set  $\{T_w c^\alpha \mid w \in \mathfrak{S}_r\}$  for  $\alpha \in \mathbb{Z}_2^r$ . Meanwhile there exists a disjoint union

$$\{T_w c^\alpha \mid w \in \mathfrak{S}_r, \alpha \in \mathbb{Z}_2^r\} = \bigcup_{\alpha \in \mathbb{Z}_2^r} \{T_w c^\alpha \mid w \in \mathfrak{S}_r\}.$$

Then part (2) follows also from Lemma 2.3.  $\square$

An  $l$ -tuple  $(\lambda_1, \dots, \lambda_l)$  of nonnegative integers  $\lambda_i \in \mathbb{N}$  is called a *composition* of  $r$  if  $\sum_{i=1}^l \lambda_i = r$ . For a composition  $\lambda = (\lambda_1, \dots, \lambda_l)$  of  $r$ , denote by  $\mathfrak{S}_\lambda = \mathfrak{S}_{\lambda_1} \times \dots \times \mathfrak{S}_{\lambda_l}$  the standard Young subgroup of  $\mathfrak{S}_r$  and let  $\mathcal{D}_\lambda$  be the set of minimal length right coset representatives of  $\mathfrak{S}_\lambda$  in  $\mathfrak{S}_r$ . Let  $\mathcal{H}_{\lambda,R}$  and  $\mathcal{H}_{\lambda,R}^c$  be the associated subalgebras of  $\mathcal{H}_{r,R}$  and  $\mathcal{H}_{r,R}^c$ , respectively. Then clearly  $\mathcal{H}_{\lambda,R}^c \cong \mathcal{H}_{\lambda,R} \otimes \mathcal{C}_r$  as  $R$ -supermodules and moreover we have algebra isomorphisms

$$\begin{aligned} \mathcal{H}_{\lambda,R} &\cong \mathcal{H}_{\lambda_1,R} \otimes \dots \otimes \mathcal{H}_{\lambda_l,R}, \\ \mathcal{H}_{\lambda,R}^c &\cong \mathcal{H}_{\lambda_1,R}^c \otimes \dots \otimes \mathcal{H}_{\lambda_l,R}^c. \end{aligned} \quad (2.4.1)$$

We remind the reader that, for two superalgebras  $\mathcal{S}$  and  $\mathcal{T}$  over  $R$ , the (super) tensor product  $\mathcal{S} \otimes \mathcal{T}$  is naturally a superalgebra, with multiplication defined by

$$(s \otimes t)(s' \otimes t') = (-1)^{\widehat{t} \widehat{s'}}(ss') \otimes (tt') \quad \text{for all } s, s' \in \mathcal{S}, t, t' \in \mathcal{T},$$

where  $t, s'$  are homogeneous. Here we used the notation  $\widehat{a} = 0$  if  $a$  is even and  $\widehat{a} = 1$  otherwise for a homogeneous element  $a$  in a superspace.

Denote by  $\Lambda(n, r) \subset \mathbb{N}^n$  the set of compositions of  $r$  with  $n$  parts. Given  $\lambda \in \Lambda(n, r)$ , set

$$x_\lambda = \sum_{w \in \mathfrak{S}_\lambda} T_w, \quad y_\lambda = \sum_{w \in \mathfrak{S}_\lambda} (-q^{-1})^{\ell(w)} T_w, \quad (2.4.2)$$

where  $\ell(w)$  is the length of  $w$ . As a super analog of the  $q$ -Schur algebra (cf. [7, 2.9]), we introduce the  $\mathbb{Q}$ - $q$ -Schur superalgebra (or the  $q$ -Schur superalgebra of type  $\mathbb{Q}$ )

$$\mathcal{Q}_q(n, r; R) = \text{End}_{\mathcal{H}_{r,R}^c} \left( \bigoplus_{\lambda \in \Lambda(n, r)} x_\lambda \mathcal{H}_{r,R}^c \right). \quad (2.4.3)$$

In particular, we write

$$\mathcal{Q}_q(n, r) := \mathcal{Q}_q(n, r; \mathcal{A}) \text{ and } \mathcal{Q}_v(n, r) := \mathcal{Q}_q(n, r; \mathcal{Z}), \quad (2.4.4)$$

where  $q = v^2$ .

Note that  $\bigoplus_{\lambda \in \Lambda(n, r)} x_\lambda \mathcal{H}_{r,R}$  is a direct sum of  $q$ -permutation modules, which is known as the tensor space used in defining  $q$ -Schur algebras. Since  $x_\lambda \mathcal{H}_{r,R}^c \cong x_\lambda \mathcal{H}_{r,R} \otimes_{\mathcal{H}_{r,R}} \mathcal{H}_{r,R}^c$  as  $\mathcal{H}_{r,R}^c$ -supermodules, which will be called a  $q$ -permutation supermodule, it is clear we have  $\mathcal{H}_{r,R}^c$ -module isomorphism

$$\bigoplus_{\lambda \in \Lambda(n, r)} x_\lambda \mathcal{H}_{r,R}^c \cong \left( \bigoplus_{\lambda \in \Lambda(n, r)} x_\lambda \mathcal{H}_{r,R} \right) \otimes_{\mathcal{H}_{r,R}} \mathcal{H}_{r,R}^c, \quad (2.4.5)$$

which is an induced  $\mathcal{H}_{r,R}^c$ -module from the tensor space. We will see in §6 that this module itself can be regarded as a tensor product of an  $R$ -supermodule.

Given a right  $\mathcal{H}_{r,R}^c$ -supermodule  $M$ , we can twist the action with  $\varphi$  to get a new  $\mathcal{H}_{r,R}^c$ -supermodule  $M^\varphi$ . Thus, as an  $R$ -module  $M^\varphi = M$ , the new  $\mathcal{H}_{r,R}^c$ -action is defined by

$$m \cdot h = m\varphi(h), \quad \text{for all } h \in \mathcal{H}_{r,R}^c, m \in M.$$

Clearly  $\mathcal{H}_{r,R}$  is stable under the involution  $\varphi$ . If  $M$  is a  $\mathcal{H}_{r,R}$ -module, we can define  $M^\varphi$  similarly. It is known that

$$(x_\lambda \mathcal{H}_{r,R})^\varphi \cong y_{\lambda^\circ} \mathcal{H}_{r,R} \cong y_\lambda \mathcal{H}_{r,R}$$

(see [7, (2.1)], or [5, Lemma 7.39]), where  $\lambda^\circ = (\lambda_l, \lambda_{l-1}, \dots, \lambda_1)$  if  $\lambda = (\lambda_1, \dots, \lambda_l)$ . This implies  $(x_\lambda \mathcal{H}_{r,R}^c)^\varphi \cong y_\lambda \mathcal{H}_{r,R}^c$  due to the facts  $x_\lambda \mathcal{H}_{r,R}^c \cong x_\lambda \mathcal{H}_{r,R} \otimes_{\mathcal{H}_{r,R}} \mathcal{H}_{r,R}^c$  and  $y_\lambda \mathcal{H}_{r,R}^c \cong y_\lambda \mathcal{H}_{r,R} \otimes_{\mathcal{H}_{r,R}} \mathcal{H}_{r,R}^c$ . Hence, we obtain a super analog of [7, Theorem 2.24(i),(ii)]

$$\mathcal{Q}_q(n, r; R) \cong \text{End}_{\mathcal{H}_{r,R}^c} \left( \bigoplus_{\lambda \in \Lambda(n, r)} y_\lambda \mathcal{H}_{r,R}^c \right).$$

**Remark 2.5.** We may also use the anti-involution  $\tau$  in Lemma 2.2(3) to turn a left  $\mathcal{H}_{r,R}^c$ -supermodule to a right  $\mathcal{H}_{r,R}^c$ -supermodule or vice versa. More precisely, if  $M$  is a finite dimensional left  $\mathcal{H}_{r,R}^c$ -supermodule, then we define a right  $\mathcal{H}_{r,R}^c$ -supermodule  $M^\tau$  by setting  $M^\tau = M$  as a  $R$ -module and making  $\mathcal{H}_{r,R}^c$ -action through  $\tau$ :

$$m \cdot h = \tau(h)m, \quad \forall h \in \mathcal{H}_{r,R}^c, m \in M.$$

In particular, if  $M = \mathcal{H}_{r,R}^c y_\lambda$ , then

$$(\mathcal{H}_{r,R}^c y_\lambda)^\tau \cong \tau(\mathcal{H}_{r,R}^c y_\lambda) = \tau(y_\lambda) \mathcal{H}_{r,R}^c = x_\lambda \mathcal{H}_{r,R}^c \quad (2.5.1)$$

by [7, (2.1)] again.

### 3. $q$ -PERMUTATION SUPERMODULES FOR $\mathcal{H}_{r,R}^c$

The  $q$ -permutation supermodules  $x_\lambda \mathcal{H}_{r,R}^c$  share certain nice properties with  $q$ -permutations modules of the Hecke algebra  $\mathcal{H}_{r,R}$  as we will see below.

Recall, for a composition  $\lambda$  of  $r$ ,  $\mathcal{D}_\lambda$  is the set of minimal length right coset representatives of  $\mathfrak{S}_\lambda$  in  $\mathfrak{S}_r$ .

**Lemma 3.1.** *Let  $\lambda$  be a composition of  $r$ . Then:–*

(1) *the right  $\mathcal{H}_{r,R}^c$ -supermodule  $x_\lambda \mathcal{H}_{r,R}^c$  (resp.,  $y_\lambda \mathcal{H}_{r,R}^c$ ) is  $R$ -free with basis*

$$\{x_\lambda T_d c^\alpha \mid d \in \mathcal{D}_\lambda, \alpha \in \mathbb{Z}_2^r\} \quad \text{resp.,} \quad \{y_\lambda T_d c^\alpha \mid d \in \mathcal{D}_\lambda, \alpha \in \mathbb{Z}_2^r\};$$

(2)  $x_\lambda \mathcal{H}_{r,R}^c = \{h \in \mathcal{H}_{r,R}^c \mid T_{s_k} h = qh, \forall s_k \in \mathfrak{S}_\lambda\}$ .

(3)  $y_\lambda \mathcal{H}_{r,R}^c = \{h \in \mathcal{H}_{r,R}^c \mid T_{s_k} h = -h, \forall s_k \in \mathfrak{S}_\lambda\}$ .

*Similar statements hold for the left  $\mathcal{H}_{r,R}^c$ -supermodules  $\mathcal{H}_{r,R}^c x_\lambda$  and  $\mathcal{H}_{r,R}^c y_\lambda$ .*

*Proof.* Part (1) is clear. We now prove part (2). Since  $T_{s_k} x_\lambda = q x_\lambda$  for all  $s_k \in \mathfrak{S}_\lambda$ , we obtain  $x_\lambda \mathcal{H}_{r,R}^c \subseteq \{h \in \mathcal{H}_{r,R}^c \mid T_{s_k} h = qh, \forall s_k \in \mathfrak{S}_\lambda\} =: \mathcal{E}$ .

Conversely, given  $h = \sum_{w \in \mathfrak{S}_r, \alpha \in \mathbb{Z}_2^r} h_{w,\alpha} T_w c^\alpha \in \mathcal{E}$  with  $h_{w,\alpha} \in R$ , write

$$h_\alpha = \sum_{w \in \mathfrak{S}_r} h_{w,\alpha} T_w c^\alpha = \left( \sum_{w \in \mathfrak{S}_r} h_{w,\alpha} T_w \right) c^\alpha.$$

Then  $h_\alpha \in \mathcal{H}_{r,R} c^\alpha$  and  $h$  can be written as  $h = \sum_{\alpha \in \mathbb{Z}_2^r} h_\alpha$ . Assume  $T_{s_k} h = qh$  for all  $s_k \in \mathfrak{S}_\lambda$ . Then we obtain

$$\sum_{\alpha \in \mathbb{Z}_2^r} (T_{s_k} h_\alpha - qh_\alpha) = 0$$

for  $s_k \in \mathfrak{S}_\lambda$ . Observe that  $T_{s_k} h_\alpha - qh_\alpha \in \mathcal{H}_{r,R} c^\alpha$  for each  $\alpha \in \mathbb{Z}_2^r$ . Then, by Corollary 2.4(2), one may deduce that  $T_{s_k} h_\alpha = qh_\alpha$  for all  $s_k \in \mathfrak{S}_\lambda$  and  $\alpha \in \mathbb{Z}_2^r$ . This implies  $T_{s_k} (\sum_{w \in \mathfrak{S}_r} h_{w,\alpha} T_w) = q (\sum_{w \in \mathfrak{S}_r} h_{w,\alpha} T_w)$  for all  $s_k \in \mathfrak{S}_\lambda$  since  $c^\alpha$  is invertible. Then using the classical result for  $\mathcal{H}_{r,R}$ , we obtain  $\sum_{w \in \mathfrak{S}_r} h_{w,\alpha} T_w \in x_\lambda \mathcal{H}_{r,R}$ . This means  $h_\alpha \in x_\lambda \mathcal{H}_{r,R}^c$  for  $\alpha \in \mathbb{Z}_2^r$ . Therefore, part (2) is proved. The proof of (3) is similar.

Applying the anti-involution  $\tau$  in Lemma 2.2(3) to (1)–(3) gives the last assertion.  $\square$

**Lemma 3.2.** *The following isomorphism of  $R$ -modules holds for compositions  $\lambda, \mu$  of  $r$ :*

$$\text{Hom}_{\mathcal{H}_{r,R}^c}(x_\mu \mathcal{H}_{r,R}^c, x_\lambda \mathcal{H}_{r,R}^c) \cong x_\lambda \mathcal{H}_{r,R}^c \cap \mathcal{H}_{r,R}^c x_\mu.$$

*Proof.* Given  $f \in \text{Hom}_{\mathcal{H}_{r,R}^c}(x_\mu \mathcal{H}_{r,R}^c, x_\lambda \mathcal{H}_{r,R}^c)$ , we have  $f(x_\mu T_{s_k}) = f(x_\mu T_{s_k}) = f(qx_\mu) = qf(x_\mu)$  for all  $s_k \in \mathfrak{S}_\mu$ . By Lemma 3.1, one can deduce that  $f(x_\mu) \in \mathcal{H}_{r,R}^c x_\mu$ . Clearly,  $f(x_\mu) \in x_\lambda \mathcal{H}_{r,R}^c$ . Therefore, we obtain an  $R$ -linear map

$$\begin{aligned} \Phi : \text{Hom}_{\mathcal{H}_{r,R}^c}(x_\mu \mathcal{H}_{r,R}^c, x_\lambda \mathcal{H}_{r,R}^c) &\longrightarrow x_\lambda \mathcal{H}_{r,R}^c \cap \mathcal{H}_{r,R}^c x_\mu, \\ f &\longmapsto f(x_\mu). \end{aligned}$$

Observe that  $\Phi$  is injective since each homomorphism  $f \in \text{Hom}_{\mathcal{H}_{r,R}^c}(x_\mu \mathcal{H}_{r,R}^c, x_\lambda \mathcal{H}_{r,R}^c)$  is uniquely determined by  $f(x_\mu)$ . On the other hand, taking an element  $z \in x_\lambda \mathcal{H}_{r,R}^c \cap \mathcal{H}_{r,R}^c x_\mu$ , we define  $f_z \in \text{Hom}_{\mathcal{H}_{r,R}^c}(x_\mu \mathcal{H}_{r,R}^c, x_\lambda \mathcal{H}_{r,R}^c)$  via  $f_z(x_\mu h) = zh$  for any  $h \in \mathcal{H}_{r,R}^c$ . It is straightforward to check that  $f_z$  is well-defined and  $\Phi(f_z) = z$ . This means that  $\Phi$  is surjective. Hence the lemma is verified.  $\square$

We will prove that the  $R$ -modules  $x_\lambda \mathcal{H}_{r,R}^c \cap \mathcal{H}_{r,R}^c x_\mu$  is free for the special case  $\lambda = \mu = (r)$  in §4 and for the general case in §5. We need some preparation. In the following, we will display a new basis for  $\mathcal{H}_{r,R}^c$  and  $x_\lambda \mathcal{H}_{r,R}^c$ . We start with recalling several facts for Hecke algebras  $\mathcal{H}_{r,R}$ .

Given two compositions  $\lambda, \mu$  of  $r$ , let  $\mathcal{D}_{\lambda,\mu} = \mathcal{D}_\lambda \cap \mathcal{D}_\mu^{-1}$ , where  $\mathcal{D}_\mu^{-1} = \{d^{-1} \mid d \in \mathcal{D}_\mu\}$  is the set of minimal length left coset representatives of  $\mathfrak{S}_\mu$  in  $\mathfrak{S}_r$ .

**Lemma 3.3** ([2], [6, Lemma 1.6]). *Suppose  $\lambda, \mu$  are compositions of  $r$ .*

- (1) *The set  $\mathcal{D}_{\lambda,\mu}$  is a system of  $\mathfrak{S}_\lambda$ - $\mathfrak{S}_\mu$  double coset representatives in  $\mathfrak{S}_r$ .*
- (2) *The element  $d \in \mathcal{D}_{\lambda,\mu}$  is the unique element of minimal length in  $\mathfrak{S}_\lambda d \mathfrak{S}_\mu$ .*
- (3) *If  $d \in \mathcal{D}_{\lambda,\mu}$ , there exists a composition  $\nu(d)$  of  $r$  such that  $\mathfrak{S}_{\nu(d)} = d^{-1} \mathfrak{S}_\lambda d \cap \mathfrak{S}_\mu$ .*
- (4) *Each element  $w \in \mathfrak{S}_\lambda d \mathfrak{S}_\mu$  can be written uniquely as  $w = u d \sigma$  with  $u \in \mathfrak{S}_\lambda$  and  $\sigma \in \mathcal{D}_{\nu(d)} \cap \mathfrak{S}_\mu$ . Moreover,  $\ell(w) = \ell(u) + \ell(d) + \ell(\sigma)$ .*

Suppose  $\lambda, \mu$  are compositions of  $r$ . Observe that  $\mathcal{D}_{\mu,\lambda} = \mathcal{D}_\mu \cap \mathcal{D}_\lambda^{-1} = \mathcal{D}_{\lambda,\mu}^{-1}$  is a system of  $\mathfrak{S}_\mu$ - $\mathfrak{S}_\lambda$  double coset representatives in  $\mathfrak{S}_r$  and hence  $d \mathfrak{S}_\mu d^{-1} \cap \mathfrak{S}_\lambda$  is a standard Young subgroup of  $\mathfrak{S}_\lambda$  for each  $d \in \mathcal{D}_{\lambda,\mu}$  by Lemma 3.3(3). Thus  $\nu(d^{-1})$  is well-defined via  $\mathfrak{S}_{\nu(d^{-1})} = d \mathfrak{S}_\mu d^{-1} \cap \mathfrak{S}_\lambda$  and moreover  $\mathfrak{S}_{\nu(d^{-1})} = d \mathfrak{S}_{\nu(d)} d^{-1}$ .

**Corollary 3.4.** *Suppose  $\lambda, \mu$  are compositions of  $r$  and  $d \in \mathcal{D}_{\lambda,\mu}$ .*

- (1) *For each  $u \in \mathfrak{S}_\lambda$ , the element  $ud$  can be written uniquely as  $ud = u' d \tau$  with  $u' \in \mathcal{D}_{\nu(d^{-1})}^{-1} \cap \mathfrak{S}_\lambda$  and  $\tau \in \mathfrak{S}_{\nu(d)}$ . Moreover  $\ell(ud) = \ell(u) + \ell(d) = \ell(u') + \ell(d) + \ell(\tau)$ .*
- (2) *The coset  $\mathfrak{S}_\lambda d = \{u' d \tau \mid u' \in \mathcal{D}_{\nu(d^{-1})}^{-1} \cap \mathfrak{S}_\lambda, \tau \in \mathfrak{S}_{\nu(d)}\}$ .*
- (3) *Each element  $w \in \mathfrak{S}_\lambda d \mathfrak{S}_\mu$  can be written uniquely as  $w = u' d \pi$  with  $u' \in \mathcal{D}_{\nu(d^{-1})}^{-1} \cap \mathfrak{S}_\lambda$  and  $\pi \in \mathfrak{S}_\mu$ .*
- (4) *For  $d_1, d_2 \in \mathcal{D}_{\lambda,\mu}$  and  $u_1 \in \mathcal{D}_{\nu(d_1^{-1})}^{-1} \cap \mathfrak{S}_\lambda, u_2 \in \mathcal{D}_{\nu(d_2^{-1})}^{-1} \cap \mathfrak{S}_\lambda$ , if  $u_1 \neq u_2$  or  $d_1 \neq d_2$ , then  $u_1 d_1 \mathfrak{S}_\mu \cap u_2 d_2 \mathfrak{S}_\mu = \emptyset$ .*

**Corollary 3.5.** *Suppose  $\lambda, \mu$  are compositions of  $r$  and  $d \in \mathcal{D}_{\lambda,\mu}$ . Then*

$$x_\lambda T_d = \left( \sum_{u' \in \mathcal{D}_{\nu(d^{-1})}^{-1} \cap \mathfrak{S}_\lambda} T_{u'} \right) T_d x_{\nu(d)}.$$

It is known that  $\mathcal{H}_{r,R}$  has the basis  $\{T_w \mid w \in \mathfrak{S}_r\}$  satisfying the following:

$$T_w T_{s_i} = \begin{cases} T_{ws_i}, & \text{if } \ell(ws_i) = \ell(w) + 1, \\ (q-1)T_w + qT_{ws_i}, & \text{if } \ell(ws_i) = \ell(w) - 1. \end{cases} \quad (3.5.1)$$

Recall that  $\leq$  is the Bruhat order of  $\mathfrak{S}_r$ . Then by (3.5.1) one can deduce that, for any  $y, w \in \mathfrak{S}_r$ , there is an element  $y * w \in \mathfrak{S}_r$  such that  $\ell(y * w) \leq \ell(y) + \ell(w)$  and

$$T_y T_w = \sum_{z \leq y * w} f_z^{y,w} T_z \quad (3.5.2)$$

for some  $f_z^{y,w} \in R$  with  $f_{y * w}^{y,w} \neq 0$  (see, e.g., [5, Proposition 4.30]).

**Lemma 3.6.** *Suppose  $\lambda, \mu \in \Lambda(n, r)$ . Let  $u \in \mathfrak{S}_\lambda, d \in \mathcal{D}_{\lambda,\mu}$ , and  $w \in \mathfrak{S}_\mu$ . Then we have*

$$T_u T_d T_w = \sum_{u' \leq u d * w} f_{u'}^{u d, w} T_{u'} \text{ for some } f_{u'}^{u d, w} \in R.$$

*In particular,  $\ell(u') \leq \ell(u) + \ell(d) + \ell(w)$  whenever  $f_{u'}^{u d, w} \neq 0$ .*

*Proof.* Clearly from Lemma 3.3(3) and (3.7.1), we have  $T_u T_d = T_{ud}$ . Then the lemma follows from (3.5.2).  $\square$

**Lemma 3.7.** *For any given  $\lambda, \mu \in \Lambda(n, r)$ , the set*

$$\mathcal{B}' = \{T_u T_d c^\alpha T_\sigma \mid u \in \mathfrak{S}_\lambda, d \in \mathcal{D}_{\lambda, \mu}, \sigma \in \mathcal{D}_{\nu(d)} \cap \mathfrak{S}_\mu, \alpha \in \mathbb{Z}_2^r\}$$

*forms an  $R$ -basis for  $\mathcal{H}_{r, R}^c$ .*

*Proof.* Let  $M$  be the  $R$ -submodule of  $\mathcal{H}_{r, R}^c$  spanned by the set  $\mathcal{B}'$ . Take an arbitrary  $w \in \mathfrak{S}_r$  and  $\alpha \in \mathbb{Z}_2^r$ . We now prove every  $T_w c^\alpha \in M$  by induction on  $\ell(w)$ . Clearly, if  $\ell(w) = 0$ , then  $T_w = 1$  and  $c^\alpha \in M$ . Assume now  $\ell(w) \geq 1$ . By Lemma 3.3(1) and (4), there exist unique  $d \in \mathcal{D}_{\lambda, \mu}$  and  $u \in \mathfrak{S}_\lambda, \sigma \in \mathcal{D}_{\nu(d)} \cap \mathfrak{S}_\mu$  such that  $w = ud\sigma$  and  $\ell(w) = \ell(u) + \ell(d) + \ell(\sigma)$ . Then by (3.5.1) we obtain  $T_w = T_u T_d T_\sigma$ . Hence, by (2.2.2) and Lemma 3.6, we have

$$\begin{aligned} T_w c^\alpha &= T_u T_d (T_\sigma c^\alpha) = \pm T_u T_d \left( c^{\alpha \cdot \sigma^{-1}} T_\sigma - \sum_{\pi < \sigma, \beta \in \mathbb{Z}_2^r} a_{\beta, \pi}^{\alpha \cdot \sigma^{-1}, \sigma} T_\pi c^\beta \right) \\ &= \pm T_u T_d c^{\alpha \cdot \sigma^{-1}} T_\sigma \mp \sum_{\pi < \sigma, \beta \in \mathbb{Z}_2^r} a_{\beta, \pi}^{\alpha \cdot \sigma^{-1}, \sigma} T_u T_d T_\pi c^\beta \\ &= \pm T_u T_d c^{\alpha \cdot \sigma^{-1}} T_\sigma \mp \sum_{\pi < \sigma, \beta \in \mathbb{Z}_2^r} a_{\beta, \pi}^{\alpha \cdot \sigma^{-1}, \sigma} \sum_{u' \leq ud * \pi, \beta \in \mathbb{Z}_2^r} f_{u', \beta}^{ud, \pi} T_{u'} c^\beta, \end{aligned} \quad (3.7.1)$$

where  $u'$  satisfying  $\ell(u') \leq \ell(u) + \ell(d) + \ell(\pi)$ . Since  $\pi < \sigma$ , we have  $\ell(u') \leq \ell(u) + \ell(d) + \ell(\pi) < \ell(u) + \ell(d) + \ell(\sigma) = \ell(w)$ . Then, by induction on  $\ell(w)$ , each term  $T_{u'} c^\beta$  on the right hand side of (3.7.1) belongs to  $M$  and so does  $T_w c^\alpha$  by (3.7.1). Hence,  $M = \mathcal{H}_{r, R}^c$ , or equivalently,  $\mathcal{H}_{r, R}^c$  is spanned by the set  $\mathcal{B}'$  by Lemma 2.3. Moreover, by (3.7.1), we observe that the transition matrix between  $\mathcal{B}'$  and the basis  $\mathcal{B} = \{T_w c^\alpha \mid w \in \mathfrak{S}_r, \alpha \in \mathbb{Z}_2^r\}$  is upper block triangular with respect to a linear order that refines the order  $T_w c^\alpha \leq T_\pi c^\beta \iff w \leq \pi$  for  $w, \pi \in \mathfrak{S}_r$  and  $\alpha, \beta \in \mathbb{Z}_2^r$ . Note that the diagonal blocks are permutation matrices and are labelled by  $w \in \mathfrak{S}_r$  of size  $2^r \times 2^r$ . Hence, the set  $\mathcal{B}'$  is linearly independent.  $\square$

For  $\lambda, \mu \in \Lambda(n, r)$ , let

$$X_{\lambda, \mu} = \left( \bigcup_{d \in \mathcal{D}_{\lambda, \mu}} (\{d\} \times (\mathcal{D}_{\nu(d)} \cap \mathfrak{S}_\mu)) \right) \times \mathbb{Z}_2^r.$$

**Corollary 3.8.** *For given  $\lambda, \mu \in \Lambda(n, r)$ , we may decompose  $\mathcal{H}_{r, R}^c$  into a direct sum of left  $\mathcal{H}_{\lambda, R}$ -modules or right  $\mathcal{H}_{\mu, R}^c$ -modules:*

$$\mathcal{H}_{r, R}^c = \bigoplus_{(d, \sigma, \alpha) \in X_{\lambda, \mu}} \mathcal{H}_{\lambda, R} T_d c^\alpha T_\sigma = \bigoplus_{d \in \mathcal{D}_{\lambda, \mu}} \bigoplus_{u \in \mathcal{D}_{\nu(d-1)}^{-1} \cap \mathfrak{S}_\lambda} T_u T_d \mathcal{H}_{\mu, R}^c.$$

*In particular, the set  $\{x_\lambda T_d c^\alpha T_\sigma \mid (d, \sigma, \alpha) \in X_{\lambda, \mu}\}$  forms a basis for  $x_\lambda \mathcal{H}_{r, R}^c$ .*

*Proof.* Clearly,  $\mathcal{H}_{\lambda, R} T_d c^\alpha T_\sigma$  is spanned by  $\{T_u T_d c^\alpha T_\sigma \mid u \in \mathfrak{S}_\lambda\}$ . Since

$$\{T_u T_d c^\alpha T_\sigma \mid u \in \mathfrak{S}_\lambda, (d, \sigma, \alpha) \in X_{\lambda, \mu}\} = \bigcup_{(d, \sigma, \alpha) \in X_{\lambda, \mu}} \{T_u T_d c^\alpha T_\sigma \mid u \in \mathfrak{S}_\lambda\},$$

a disjoint union, the assertion follows from Lemma 3.7 and the Corollary 3.4.  $\square$

4. THE ELEMENTS  $c_{q,i,j}$ 

For  $r \geq 1$  and  $1 \leq i < j \leq r$ , we set

$$c_{q,i,j} = q^{j-i}c_i + q^{j-i-1}c_{i+1} + \cdots + qc_{j-1} + c_j, \quad c'_{q,i,j} = c_i + qc_{i+1} + \cdots + q^{j-i}c_j \quad (4.0.1)$$

For simplicity, we write  $c_{q,r} = c_{q,1,r}$ . Then we have

$$c_{q,r} = q^{r-1}c_1 + q^{r-2}c_2 + \cdots + c_r, \quad c'_{q,r} = c_1 + qc_2 + \cdots + q^{r-1}c_r. \quad (4.0.2)$$

**Lemma 4.1.** *Let  $r \geq 1$ . The following holds in  $\mathcal{H}_{r,R}^c$ :*

$$x_{(r)}c_{q,r} = c'_{q,r}x_{(r)} \quad \text{and} \quad y_{(r)}c'_{q,r} = c_{q,r}y_{(r)}.$$

*Proof.* We only prove the first formula. The proof of the second one is similar. We apply induction on  $r$ . It is known that

$$x_{(r)} = x_{(r-1)}(1 + T_{r-1} + T_{r-1}T_{r-2} + \cdots + T_{r-1} \cdots T_1) \quad (4.1.1)$$

since  $\{1, s_{r-1}, s_{r-1}s_{r-2}, \dots, s_{r-1} \cdots s_2s_1\} = \mathcal{D}_{(r-1,1)}$ . Using (2.0.2), a direct calculation shows for  $1 \leq k \leq r-1$  and  $1 \leq l \leq r$ :

$$T_{r-1} \cdots T_k c_l = \begin{cases} c_l T_{r-1} \cdots T_k, & \text{if } 1 \leq l \leq k-1, \\ c_r T_{r-1} \cdots T_k, & \text{if } l = k, \\ c_{l-1} T_{r-1} \cdots T_k - (q-1)T_{l-2} \cdots T_k (c_k - c_r) T_{r-1} \cdots T_l, & \text{if } k+1 \leq l \leq r, \end{cases}$$

where we use the convention  $T_{r-1} \cdots T_l = 1$  if  $l = r$  and the formula in the last case is due to the following computation:

$$\begin{aligned} T_{r-1} \cdots T_k c_l &= T_{r-1} \cdots T_l \cdot (c_{l-1} T_{l-1} - (q-1)(c_{l-1} - c_l)) \cdot T_{l-2} \cdots T_k \\ &= c_{l-1} T_{r-1} \cdots T_l T_{l-1} T_{l-2} \cdots T_k - (q-1)(T_{r-1} \cdots T_l) \cdot (c_{l-1} - c_l) \cdot (T_{l-2} \cdots T_k) \\ &= c_{l-1} T_{r-1} \cdots T_k - (q-1)(c_{l-1} - c_r) \cdot (T_{r-1} \cdots T_l) \cdot (T_{l-2} \cdots T_k) \\ &= c_{l-1} T_{r-1} \cdots T_k - (q-1)(c_{l-1} - c_r) \cdot (T_{l-2} \cdots T_k) \cdot (T_{r-1} \cdots T_l) \\ &= c_{l-1} T_{r-1} \cdots T_k - (q-1)(T_{l-2} \cdots T_k) \cdot (c_k - c_r) \cdot (T_{r-1} \cdots T_l). \end{aligned}$$

This implies that the following holds for  $1 \leq k \leq r-1$  and  $1 \leq l \leq r$ :

$$\begin{aligned} &x_{(r-1)} T_{r-1} \cdots T_k c_l \\ &= \begin{cases} x_{(r-1)} c_l T_{r-1} \cdots T_k, & \text{if } 1 \leq l \leq k-1, \\ x_{(r-1)} c_r T_{r-1} \cdots T_k, & \text{if } l = k, \\ x_{(r-1)} c_{l-1} T_{r-1} \cdots T_k - q^{l-k-1}(q-1)x_{(r-1)}(c_k - c_r) T_{r-1} \cdots T_l, & \text{if } k+1 \leq l \leq r, \end{cases} \end{aligned}$$

where the formula in the last case is due to the fact  $x_{(r-1)} T_{l-2} \cdots T_k = q^{l-k-1} x_{(r-1)}$ . Hence we obtain

$$\begin{aligned} x_{(r-1)} T_{r-1} \cdots T_k c_{q,r} &= x_{(r-1)} T_{r-1} \cdots T_k (q^{r-k+1} c_{q,1,k-1} + q^{r-k} c_k + c_{q,k+1,r}) \\ &= x_{(r-1)} (q^{r-k+1} c_{q,1,k-1} + q^{r-k} c_r + c_{q,k,r-1}) T_{r-1} \cdots T_k \\ &\quad - \sum_{l=k+1}^r q^{r-l} q^{l-k-1} (q-1) x_{(r-1)} (c_k - c_r) T_{r-1} \cdots T_l \quad (\text{noting } T_{r-1} \cdots T_l = 1 \text{ if } l = r) \\ &= x_{(r-1)} (q^{r-k+1} c_{q,1,k-1} + q^{r-k} c_r + c_{q,k,r-1}) T_{r-1} \cdots T_k \\ &\quad - \sum_{l=k+1}^{r-1} (q-1) q^{r-k-1} x_{(r-1)} (c_k - c_r) T_{r-1} \cdots T_l - (q-1) q^{r-k-1} x_{(r-1)} (c_k - c_r). \end{aligned}$$

This together with (4.1.1) gives rise to

$$\begin{aligned}
x_{(r)}c_{q,r} &= (x_{(r-1)} + \sum_{k=1}^{r-1} x_{(r-1)}T_{r-1} \cdots T_k)c_{q,r} \\
&= x_{(r-1)}c_{q,r} + \sum_{k=1}^{r-1} x_{(r-1)}(q^{r-k+1}c_{q,1,k-1} + q^{r-k}c_r + c_{q,k,r-1})T_{r-1} \cdots T_k \\
&\quad - \sum_{k=1}^{r-1} \sum_{l=k+1}^{r-1} (q-1)q^{r-k-1}x_{(r-1)}(c_k - c_r)T_{r-1} \cdots T_l \\
&\quad - \sum_{k=1}^{r-1} (q-1)q^{r-k-1}x_{(r-1)}(c_k - c_r).
\end{aligned} \tag{4.1.2}$$

Combining the first term and the last sum in (4.1.2) yields

$$\begin{aligned}
x_{(r-1)}c_{q,r} - \sum_{k=1}^{r-1} x_{(r-1)}(q-1)q^{r-k-1}(c_k - c_r) \\
&= x_{(r-1)} \left( \sum_{k=1}^r q^{r-k}c_k - \sum_{k=1}^{r-1} (q^{r-k} - q^{r-k-1})c_k + \sum_{k=1}^{r-1} (q^{r-k} - q^{r-k-1})c_r \right) \\
&= x_{(r-1)}(q^{r-2}c_1 + q^{r-3}c_2 + \cdots + c_{r-1} + q^{r-1}c_r) = x_{(r-1)}(c_{q,1,r-1} + q^{r-1}c_r).
\end{aligned}$$

Also, the second sum in (4.1.2) can be rewritten as

$$\begin{aligned}
x_{(r-1)} \sum_{k=1}^{r-1} \sum_{l=k+1}^{r-1} (q-1)q^{r-k-1}(c_k - c_r)T_{r-1} \cdots T_l \\
&= x_{(r-1)} \sum_{l=2}^{r-1} \left( \sum_{k=1}^{l-1} (q-1)q^{r-k-1}(c_k - c_r) \right) T_{r-1} \cdots T_l \\
&= x_{(r-1)} \sum_{l=2}^{r-1} ((q^{r-l+1} - q^{r-l})c_{q,1,l-1} - (q^{r-1} - q^{r-l})c_r) T_{r-1} \cdots T_l.
\end{aligned}$$

Hence, substituting into (4.1.2) yields

$$x_{(r)}c_{q,r} = x_{(r-1)}(c_{q,1,r-1} + q^{r-1}c_r)(1 + \sum_{k=1}^{r-1} T_{r-1} \cdots T_k). \tag{4.1.3}$$

Here the following calculation has been used for  $2 \leq k \leq r-1$ :

$$\begin{aligned}
&(q^{r-k+1}c_{q,1,k-1} + q^{r-k}c_r + c_{q,k,r-1}) - ((q^{r-k+1} - q^{r-k})c_{q,1,k-1} - (q^{r-1} - q^{r-k})c_r) \\
&= q^{r-k}c_{q,1,k-1} + c_{q,k,r-1} + q^{r-1}c_r = c_{q,1,r-1} + q^{r-1}c_r = c_{q,r-1} + q^{r-1}c_r,
\end{aligned}$$

which is also the first ( $k=1$ ) term of the first sum in (4.1.2). By induction on  $r$ , we have

$$x_{(r-1)}c_{q,1,r-1} = c'_{q,r-1}x_{(r-1)}$$

and hence, by (4.0.2), one can deduce

$$x_{(r-1)}(c_{q,1,r-1} + q^{r-1}c_r) = (c'_{q,r-1} + q^{r-1}c_r)x_{(r-1)} = c'_{q,r}x_{(r-1)}.$$

Thus, by (4.1.3) and (4.1.1), we obtain

$$x_{(r)}c_{q,r} = c'_{q,r}x_{(r-1)}(1 + T_{r-1} + \cdots + T_{r-1} \cdots T_1) = c'_{q,r}x_{(r)}.$$

This proves the lemma.  $\square$

**Proposition 4.2.** *For  $r \geq 1$ , the intersection  $x_{(r)}\mathcal{H}_{r,R}^c \cap \mathcal{H}_{r,R}^c x_{(r)}$  is a free  $R$ -module with basis  $\{x_{(r)}, x_{(r)}c_{q,r}\}$ .*

*Proof.* By Lemma 4.1, the set  $\{x_{(r)}, x_{(r)}c_{q,r}\}$  is contained in the intersection  $x_{(r)}\mathcal{H}_{r,R}^c \cap \mathcal{H}_{r,R}^c x_{(r)}$  and clearly it is linearly independent. It remains to show that  $x_{(r)}\mathcal{H}_{r,R}^c \cap \mathcal{H}_{r,R}^c x_{(r)}$  is spanned by  $\{x_{(r)}, x_{(r)}c_{q,r}\}$ . Take an arbitrary  $z \in x_{(r)}\mathcal{H}_{r,R}^c \cap \mathcal{H}_{r,R}^c x_{(r)}$ . Since  $z \in x_{(r)}\mathcal{H}_{r,R}^c$ , by Lemma 3.1(1), we can write  $z$  as  $z = \sum_{\alpha \in \mathbb{Z}_2^r} f_{\alpha} x_{(r)} c^{\alpha}$ , with  $f_{\alpha} \in R$ . Moreover since  $z \in \mathcal{H}_{r,R}^c x_{(r)}$ , by Lemma 3.1(2) we have

$$zT_{s_k} = qz, \quad \forall 1 \leq k \leq r-1. \quad (4.2.1)$$

For  $\alpha = (\alpha_1, \dots, \alpha_r)$  and  $1 \leq k \leq r-1$ , by (2.0.2) and (2.2.1), we have

$$\begin{aligned} c_j^{\alpha_j} T_{s_k} &= T_{s_k} c_j^{\alpha_j}, \quad \text{for } j \neq k, k+1, \\ c_{k+1}^{\alpha_{k+1}} T_{s_k} &= T_{s_k} c_k^{\alpha_{k+1}}, \quad c_k^{\alpha_k} T_{s_k} = T_{s_k} c_{k+1}^{\alpha_k} + (q-1)(c_k^{\alpha_k} - c_{k+1}^{\alpha_k}), \\ c_k^{\alpha_k} c_{k+1}^{\alpha_{k+1}} T_{s_k} &= T_{s_k} c_{k+1}^{\alpha_k} c_k^{\alpha_{k+1}} + (q-1)(c_k^{\alpha_k + \alpha_{k+1}} - c_{k+1}^{\alpha_k} c_k^{\alpha_{k+1}}). \end{aligned}$$

Then, by the fact  $x_{(r)}T_{s_k} = qx_{(r)}$  for  $1 \leq k \leq r-1$ , we obtain

$$\begin{aligned} x_{(r)}c^{\alpha} T_{s_k} &= x_{(r)}T_{s_k} c_1^{\alpha_1} \cdots c_{k-1}^{\alpha_{k-1}} c_{k+1}^{\alpha_{k+1}} c_k^{\alpha_k} c_{k+2}^{\alpha_{k+2}} \cdots c_r^{\alpha_r} \\ &\quad + (q-1)x_{(r)}c_1^{\alpha_1} \cdots c_{k-1}^{\alpha_{k-1}} (c_k^{\alpha_k + \alpha_{k+1}} - c_{k+1}^{\alpha_k} c_k^{\alpha_{k+1}}) c_{k+2}^{\alpha_{k+2}} \cdots c_r^{\alpha_r} \\ &= x_{(r)}c_1^{\alpha_1} \cdots c_{k-1}^{\alpha_{k-1}} c_{k+1}^{\alpha_k} c_k^{\alpha_{k+1}} c_{k+2}^{\alpha_{k+2}} \cdots c_r^{\alpha_r} \\ &\quad + (q-1)x_{(r)}c_1^{\alpha_1} \cdots c_{k-1}^{\alpha_{k-1}} c_k^{\alpha_k + \alpha_{k+1}} c_{k+2}^{\alpha_{k+2}} \cdots c_r^{\alpha_r}. \end{aligned} \quad (4.2.2)$$

This together with (4.2.1) implies

$$\begin{aligned} q \sum_{\alpha \in \mathbb{Z}_2^r} f_{\alpha} x_{(r)} c^{\alpha} &= zT_{s_k} = \sum_{\alpha \in \mathbb{Z}_2^r} f_{\alpha} x_{(r)} c_1^{\alpha_1} \cdots c_{k-1}^{\alpha_{k-1}} c_{k+1}^{\alpha_{k+1}} c_k^{\alpha_k} c_{k+2}^{\alpha_{k+2}} \cdots c_r^{\alpha_r} \\ &\quad + \sum_{\alpha \in \mathbb{Z}_2^r} (q-1) f_{\alpha} x_{(r)} c_1^{\alpha_1} \cdots c_{k-1}^{\alpha_{k-1}} c_k^{\alpha_k + \alpha_{k+1}} c_{k+2}^{\alpha_{k+2}} \cdots c_r^{\alpha_r} \end{aligned}$$

for each  $1 \leq k \leq r-1$ . Then by using Lemma 3.1(1) and by equating the coefficients of  $x_{(r)}c^{\alpha}$  on both sides for each  $\alpha \in \mathbb{Z}_2^r$ , we obtain

$$\begin{aligned} qf_{(\alpha_1, \dots, \alpha_{k-1}, 1, 1, \alpha_{k+2}, \dots, \alpha_r)} &= -f_{(\alpha_1, \dots, \alpha_{k-1}, 1, 1, \alpha_{k+2}, \dots, \alpha_r)}, \\ qf_{(\alpha_1, \dots, \alpha_{k-1}, 0, 1, \alpha_{k+2}, \dots, \alpha_r)} &= f_{(\alpha_1, \dots, \alpha_{k-1}, 1, 0, \alpha_{k+2}, \dots, \alpha_r)}, \\ qf_{(\alpha_1, \dots, \alpha_{k-1}, 0, 0, \alpha_{k+2}, \dots, \alpha_r)} &= f_{(\alpha_1, \dots, \alpha_{k-1}, 0, 0, \alpha_{k+2}, \dots, \alpha_r)} \\ &\quad + (q-1)(f_{(\alpha_1, \dots, \alpha_{k-1}, 1, 1, \alpha_{k+2}, \dots, \alpha_r)} + f_{(\alpha_1, \dots, \alpha_{k-1}, 0, 0, \alpha_{k+2}, \dots, \alpha_r)}), \\ qf_{(\alpha_1, \dots, \alpha_{k-1}, 1, 0, \alpha_{k+2}, \dots, \alpha_r)} &= f_{(\alpha_1, \dots, \alpha_{k-1}, 0, 1, \alpha_{k+2}, \dots, \alpha_r)} \\ &\quad + (q-1)(f_{(\alpha_1, \dots, \alpha_{k-1}, 1, 0, \alpha_{k+2}, \dots, \alpha_r)} + f_{(\alpha_1, \dots, \alpha_{k-1}, 0, 1, \alpha_{k+2}, \dots, \alpha_r)}), \end{aligned} \quad (4.2.3)$$

where the relation (2.0.1) is used in the first and last equalities. Hence, since the characteristic of  $R$  is not equal to 2, (4.2.3) implies

$$f_{(\alpha_1, \dots, \alpha_{k-1}, 1, 0, \alpha_{k+1}, \dots, \alpha_r)} = q f_{(\alpha_1, \dots, \alpha_{k-1}, 0, 1, \alpha_{k+1}, \dots, \alpha_r)}, \quad f_{(\alpha_1, \dots, \alpha_{k-1}, 1, 1, \alpha_{k+1}, \dots, \alpha_r)} = 0 \quad (4.2.4)$$

for all  $\alpha_i \in \mathbb{Z}_2$  and  $1 \leq k \leq r-1$ . Then one can deduce that

$$\begin{cases} f_{\epsilon_k} = q^{r-k} f_{\epsilon_r}, & \text{for } 1 \leq k \leq r, \\ f_{\alpha} = 0, & \text{for } \alpha \in \mathbb{Z}_2^r \setminus \mathbf{0} \text{ and } \alpha \neq \epsilon_k, 1 \leq k \leq r, \end{cases}$$

where  $\epsilon_k = (0, \dots, 0, 1, 0, \dots, 0)$  with 1 in the  $k$ -th position. This means

$$z = \sum_{\alpha \in \mathbb{Z}_2^r} f_{\alpha} x_{(r)} c^{\alpha} = f_{(0, \dots, 0)} x_{(r)} + f_{\epsilon_r} x_{(r)} (q^{r-1} c_1 + q^{r-2} c_2 + \dots + c_r) = f_{(0, \dots, 0)} x_{(r)} + f_{\epsilon_r} x_{(r)} c_{q,r}.$$

Hence  $z$  lies in the  $R$ -module spanned by  $\{x_{(r)}, x_{(r)} c_{q,r}\}$  and the proposition is verified.  $\square$

Given  $\lambda = (\lambda_1, \dots, \lambda_N) \in \Lambda(N, r)$  and  $\alpha \in \mathbb{Z}_2^N$  with  $N \geq 1$  and  $\alpha_k \leq \lambda_k$  for  $1 \leq k \leq N$ , we let  $\tilde{\lambda}_k = \lambda_1 + \dots + \lambda_k$  for  $1 \leq k \leq N$  and introduce the following elements in  $\mathcal{C}_r$ :

$$\begin{aligned} c_{\tilde{\lambda}}^{\alpha} &= (c_{q,1,\tilde{\lambda}_1})^{\alpha_1} (c_{q,\tilde{\lambda}_1+1,\tilde{\lambda}_2})^{\alpha_2} \dots (c_{q,\tilde{\lambda}_{N-1}+1,\tilde{\lambda}_N})^{\alpha_N}, \\ (c_{\tilde{\lambda}}^{\alpha})' &= (c'_{q,1,\tilde{\lambda}_1})^{\alpha_1} (c'_{q,\tilde{\lambda}_1+1,\tilde{\lambda}_2})^{\alpha_2} \dots (c'_{q,\tilde{\lambda}_{N-1}+1,\tilde{\lambda}_N})^{\alpha_N}. \end{aligned} \quad (4.2.5)$$

For notational simplicity, we define for  $\lambda, \mu \in \mathbb{N}^N$

$$\lambda \leq \mu \iff \lambda_i \leq \mu_i \text{ for all } i = 1, 2, \dots, N.$$

**Corollary 4.3.** *Suppose  $\lambda = (\lambda_1, \dots, \lambda_N) \in \Lambda(N, r)$ . Then  $x_{\lambda} \mathcal{H}_{\lambda,R}^c \cap \mathcal{H}_{\lambda,R}^c x_{\lambda}$  is a free  $R$ -module with basis  $\{x_{\lambda} c_{\tilde{\lambda}}^{\alpha} = (c_{\tilde{\lambda}}^{\alpha})' x_{\lambda} \mid \alpha \in \mathbb{Z}_2^N, \alpha \leq \lambda\}$ .*

*Proof.* Clearly, by Lemma 4.1 and (2.0.1), we have  $x_{\lambda} c_{\tilde{\lambda}}^{\alpha} = (c_{\tilde{\lambda}}^{\alpha})' x_{\lambda}$  for  $\alpha \in \mathbb{Z}_2^N$  and the set is linearly independent. We may modify the proof of Proposition 4.2 to prove the assertion. Without loss of generality, we may assume all  $\lambda_i \neq 0$ . For  $z = \sum_{\alpha \in \mathbb{Z}_2^r} f_{\alpha} x_{\lambda} c^{\alpha} \in x_{\lambda} \mathcal{H}_{\lambda,R}^c \cap \mathcal{H}_{\lambda,R}^c x_{\lambda}$ , (4.2.4) continues to hold for all  $k$  such that  $\tilde{\lambda}_{i-1} + 1 \leq k < \tilde{\lambda}_i$  for all  $i$  (assuming  $\lambda_0 = 0$ ). Then we first observe that if there exist  $\tilde{\lambda}_{i-1} + 1 \leq k < l \leq \tilde{\lambda}_i$  for some  $1 \leq i \leq N$  such that  $\alpha_k = \alpha_l = 1$ , then  $f_{\alpha} = 0$ . Now it remains to consider those  $f_{\alpha}$  for  $\alpha = \mathbf{0}$  or for  $\alpha \in \mathbb{Z}_2^r$  having the form  $\alpha = \epsilon_{k_1} + \epsilon_{k_2} + \dots + \epsilon_{k_t}$  for some  $1 \leq t \leq N$ , where there exist  $1 \leq i_1 < i_2 < \dots < i_t \leq N$  such that  $\tilde{\lambda}_{i_1-1} + 1 \leq k_1 \leq \tilde{\lambda}_{i_1}$ ,  $\tilde{\lambda}_{i_2-1} + 1 \leq k_2 \leq \tilde{\lambda}_{i_2}$ ,  $\dots$ ,  $\tilde{\lambda}_{i_t-1} + 1 \leq k_t \leq \tilde{\lambda}_{i_t}$ . In the latter case, by (4.2.4), one can deduce that

$$f_{\alpha} = q^{\tilde{\lambda}_{i_1}-k_1} q^{\tilde{\lambda}_{i_2}-k_2} \dots q^{\tilde{\lambda}_{i_t}-k_t} f_{\epsilon_{\tilde{\lambda}_{i_1}} + \epsilon_{\tilde{\lambda}_{i_2}} + \dots + \epsilon_{\tilde{\lambda}_{i_t}}}.$$

Hence,

$$z = f_{\mathbf{0}} x_{\lambda} + \sum_{t=1}^N \sum_{1 \leq i_1 < \dots < i_t \leq N} f_{\epsilon_{\tilde{\lambda}_{i_1}} + \epsilon_{\tilde{\lambda}_{i_2}} + \dots + \epsilon_{\tilde{\lambda}_{i_t}}} x_{\lambda} c_{q,\tilde{\lambda}_{i_1-1}+1,\tilde{\lambda}_{i_1}} c_{q,\tilde{\lambda}_{i_2-1}+1,\tilde{\lambda}_{i_2}} \dots c_{q,\tilde{\lambda}_{i_t-1}+1,\tilde{\lambda}_{i_t}},$$

proving the spanning condition.  $\square$

**Remark 4.4.** By (2.4.1), we have  $\mathcal{H}_{\lambda,R}^c \cong \mathcal{H}_{\lambda_1,R}^c \otimes \dots \otimes \mathcal{H}_{\lambda_N,R}^c$ , mapping  $x_{\lambda}$  to  $x_{(\lambda_1)} \otimes \dots \otimes x_{(\lambda_N)}$ . Hence, as free  $R$ -modules,

$$x_{\lambda} \mathcal{H}_{\lambda,R}^c \cong x_{(\lambda_1)} \mathcal{H}_{\lambda_1,R}^c \otimes \dots \otimes x_{(\lambda_N)} \mathcal{H}_{\lambda_N,R}^c, \quad \mathcal{H}_{\lambda,R}^c x_{\lambda} \cong \mathcal{H}_{\lambda_1,R}^c x_{(\lambda_1)} \otimes \dots \otimes \mathcal{H}_{\lambda_N,R}^c x_{(\lambda_N)}.$$

The corollary above implies the following isomorphism of free  $R$ -modules

$$x_{\lambda} \mathcal{H}_{\lambda,R}^c \cap \mathcal{H}_{\lambda,R}^c x_{\lambda} \cong (x_{(\lambda_1)} \mathcal{H}_{\lambda_1,R}^c \cap \mathcal{H}_{\lambda_1,R}^c x_{(\lambda_1)}) \otimes \dots \otimes (x_{(\lambda_N)} \mathcal{H}_{\lambda_N,R}^c \cap \mathcal{H}_{\lambda_N,R}^c x_{(\lambda_N)}).$$

## 5. INTEGRAL BASES AND THE BASE CHANGE PROPERTY

We are now ready to prove, by constructing a basis, that the module  $x_\lambda \mathcal{H}_{r,R}^c \cap \mathcal{H}_{r,R}^c x_\mu$  occurred in Lemma 3.2 is  $R$ -free. This will provide an integral  $R$ -basis for  $\mathcal{Q}_q(n, r; R)$ .

Let  $M_n(\mathbb{N})$  be the set of  $n \times n$ -matrices  $M = (m_{ij})$ . Given  $M \in M_n(\mathbb{N})$ , define

$$\text{ro}(M) = \left( \sum_j m_{1j}, \dots, \sum_j m_{nj} \right), \quad \text{co}(M) = \left( \sum_j m_{j1}, \dots, \sum_j m_{jn} \right).$$

Let

$$M_n(\mathbb{N})_r = \{M \in M_n(\mathbb{N}) \mid \sum m_{ij} = r\}.$$

Then  $\text{ro}(M), \text{co}(M) \in \Lambda(n, r)$  for any  $M \in M_n(\mathbb{N})_r$ . Given a composition  $\lambda \in \Lambda(n, r)$  and  $1 \leq k \leq n$ , define the subsets  $R_k^\lambda \subseteq \{1, \dots, r\}$  as follows:

$$R_k^\lambda = \{\tilde{\lambda}_{k-1} + 1, \tilde{\lambda}_{k-1} + 2, \dots, \tilde{\lambda}_{k-1} + \lambda_k\}.$$

It is well known that double cosets of the symmetric group can be described in terms of matrices as follows. There is a bijection

$$\begin{aligned} \mathbf{j} : M_n(\mathbb{N})_r &\longrightarrow \mathfrak{J}_n(r) = \{(\lambda, d, \mu) \mid \lambda, \mu \in \Lambda(n, r), d \in \mathcal{D}_{\lambda, \mu}\} \\ M &\longmapsto (\text{ro}(M), d_M, \text{co}(M)), \end{aligned} \quad (5.0.1)$$

where  $d_M \in \mathcal{D}_{\lambda, \mu}$  is the permutation satisfying  $|R_i^{\text{ro}(M)} \cap d_M(R_j^{\text{co}(M)})| = m_{ij}$  for  $1 \leq i, j \leq n$  with  $M = (m_{ij})$ . Moreover, the composition  $\nu(d_M)$  (resp.  $\nu(d_M^{-1})$ ), defined in Lemma 3.3(3), is obtained by reading the entries in  $M$  along columns (resp. rows), that is,

$$\begin{aligned} \nu(d_M) &= (m_{11}, \dots, m_{n1}, m_{12}, \dots, m_{n2}, \dots, m_{1n}, \dots, m_{nn}), \\ \nu(d_M^{-1}) &= (m_{11}, \dots, m_{1n}, m_{21}, \dots, m_{2n}, \dots, m_{n1}, \dots, m_{nn}). \end{aligned} \quad (5.0.2)$$

For the subset  $\mathbb{Z}_2 = \{0, 1\}$  of  $\mathbb{N}$ , let  $M_n(\mathbb{Z}_2)$  be the set of  $n \times n$ -matrices  $B = (b_{ij})$  with  $b_{ij} \in \mathbb{Z}_2$ . Set

$$\begin{aligned} M_n(\mathbb{N}|\mathbb{Z}_2) &:= \{(A|B) \mid A \in M_n(\mathbb{N}), B \in M_n(\mathbb{Z}_2)\}, \\ M_n(\mathbb{N}|\mathbb{Z}_2)_r &:= \{(A|B) \in M_n(\mathbb{N}|\mathbb{Z}_2) \mid A + B \in M_n(\mathbb{N})_r\}. \end{aligned}$$

Given  $(A|B) \in M_n(\mathbb{N}|\mathbb{Z}_2)_r$  with  $A = (a_{ij}), B = (b_{ij})$ , let  $m_{ij} = a_{ij} + b_{ij}$  for  $1 \leq i, j \leq n$  and then we have  $A + B = (m_{ij}) \in M_n(\mathbb{N})_r$ . Hence by (5.0.1), we have the permutation  $d_{A+B} \in \mathcal{D}_{\lambda, \mu}$  with  $\lambda = \text{ro}(A + B), \mu = \text{co}(A + B)$  and, moreover,

$$\nu_{A|B} := \nu(d_{A+B}) = (m_{11}, \dots, m_{n1}, m_{12}, \dots, m_{n2}, \dots, m_{1n}, \dots, m_{nn}).$$

Let  $\alpha_B = (b_{11}, \dots, b_{n1}, \dots, b_{1n}, \dots, b_{nn}) \in \mathbb{Z}_2^{n^2}$ . Then by (4.2.5) we are ready to introduce the following elements:

$$c_{A|B} = c_{\nu_{A|B}}^{\alpha_B} \in \mathcal{C}_r, \quad (5.0.3)$$

$$T_{A|B} = x_\lambda T_{d_{A+B}} c_{A|B} \sum_{\sigma \in \mathcal{D}_{\nu_{A|B}} \cap \mathfrak{S}_\mu} T_\sigma. \quad (5.0.4)$$

Given  $\lambda, \mu \in \Lambda(n, r)$  and  $d \in \mathcal{D}_{\lambda, \mu}$ , let

$$\begin{aligned} M_n(\mathbb{N}|\mathbb{Z}_2)_{\lambda, \mu} &:= \{(A|B) \in M_n(\mathbb{N}|\mathbb{Z}_2) \mid \text{ro}(A + B) = \lambda, \text{co}(A + B) = \mu\}, \\ M_n(\mathbb{N}|\mathbb{Z}_2)_{\lambda, \mu}^d &:= \{(A|B) \in M_n(\mathbb{N}|\mathbb{Z}_2)_{\lambda, \mu} \mid d_{A+B} = d\}. \end{aligned}$$

Note that, by sending  $A$  to  $(A|0)$ , we may regard  $M_n(\mathbb{N})_{\lambda,\mu} = \{A \in M_n(\mathbb{N}) \mid \text{ro}(A) = \lambda, \text{co}(A) = \mu\}$  as a subset of  $M_n(\mathbb{N}|\mathbb{Z}_2)_{\lambda,\mu}$ . Note also that

$$M_n(\mathbb{N}|\mathbb{Z}_2)_{\lambda,\mu} = \bigcup_{d \in \mathcal{D}_{\lambda,\mu}} M_n(\mathbb{N}|\mathbb{Z}_2)_{\lambda,\mu}^d. \quad (5.0.5)$$

**Lemma 5.1.** *For  $\lambda, \mu \in \Lambda(n, r)$ , the set  $\{T_{A|B} \mid (A|B) \in M_n(\mathbb{N}|\mathbb{Z}_2)_{\lambda,\mu}\}$  is a linearly independent subset of  $x_\lambda \mathcal{H}_{r,R}^c \cap \mathcal{H}_{r,R}^c x_\mu$ .*

*Proof.* We first prove that  $T_{A|B} \in x_\lambda \mathcal{H}_{r,R}^c \cap \mathcal{H}_{r,R}^c x_\mu$  for each  $(A|B) \in M_n(\mathbb{N}|\mathbb{Z}_2)_{\lambda,\mu}$ . Clearly,  $T_{A|B} \in x_\lambda \mathcal{H}_{r,R}^c$  by (5.0.4). Let  $d = d_{A+B}$ . Then by (5.0.2)

$$\nu(d) = (m_{11}, \dots, m_{n1}, m_{12}, \dots, m_{n2}, \dots, m_{1n}, \dots, m_{nn}),$$

where  $A + B = (m_{ij})$ . Hence, by Corollary 4.3 and (5.0.3),

$$x_{\nu(d)} c_{A|B} = c'_{A|B} x_{\nu(d)}, \quad (5.1.1)$$

where  $c'_{A|B} = (c_{\nu_{A|B}}^{\alpha_B})'$ . On the other hand, the partition

$$\mathfrak{S}_\mu = \dot{\bigcup}_{\sigma \in \mathcal{D}_{\nu(d)} \cap \mathfrak{S}_\mu} \mathfrak{S}_{\nu(d)\sigma} \quad (5.1.2)$$

implies

$$x_\mu = x_{\nu(d)} \sum_{\sigma \in \mathcal{D}_{\nu(d)} \cap \mathfrak{S}_\mu} T_\sigma. \quad (5.1.3)$$

Hence, by (5.0.4), Corollary 3.5, (5.1.1) and (5.1.3), one can deduce that

$$\begin{aligned} T_{A|B} &= \sum_{u' \in \mathcal{D}_{\nu(d-1)}^{-1} \cap \mathfrak{S}_\lambda} T_{u'} T_d x_{\nu(d)} c_{A|B} \sum_{\sigma \in \mathcal{D}_{\nu(d)} \cap \mathfrak{S}_\mu} T_\sigma \\ &= \sum_{u' \in \mathcal{D}_{\nu(d-1)}^{-1} \cap \mathfrak{S}_\lambda} T_{u'} T_d c'_{A|B} x_{\nu(d)} \sum_{\sigma \in \mathcal{D}_{\nu(d)} \cap \mathfrak{S}_\mu} T_\sigma = \sum_{u' \in \mathcal{D}_{\nu(d-1)}^{-1} \cap \mathfrak{S}_\lambda} T_{u'} T_d c'_{(A|B)} x_\mu. \end{aligned}$$

proving  $T_{A|B} \in \mathcal{H}_{r,R}^c x_\mu$ .

We now prove the set is linearly independent. Suppose

$$\sum_{(A|B) \in M_n(\mathbb{N}|\mathbb{Z}_2)_{\lambda,\mu}} f_{A|B} T_{A|B} = 0.$$

By the new basis  $\mathcal{B}'$  on Lemma 3.7 and (5.0.5), we have, for every  $d \in \mathcal{D}_{\lambda,\mu}$ ,

$$\sum_{(A|B) \in M_n(\mathbb{N}|\mathbb{Z}_2)_{\lambda,\mu}^d} f_{A|B} T_{A|B} = 0 \quad (5.1.4)$$

Let  $w_\lambda^0$  be the longest element in the Young subgroup  $\mathfrak{S}_\lambda$ . By (5.1.2) and the fact that  $\ell(w\sigma) = \ell(w) + \ell(\sigma)$  for  $w \in \mathfrak{S}_{\nu(d)}$  and  $\sigma \in \mathcal{D}_{\nu(d)} \cap \mathfrak{S}_\mu$ , there exists a unique longest element  $\sigma_0$  in  $\mathcal{D}_{\nu(d)} \cap \mathfrak{S}_\mu$  such that  $w_\mu^0 = w_{\nu(d)}^0 \sigma_0$ . Suppose  $(A|B) \in M_n(\mathbb{N}|\mathbb{Z}_2)_{\lambda,\mu}^d$ . Then by (2.4.2), Lemma 3.6 and (2.2.2), we obtain

$$\begin{aligned} T_{A|B} &= T_{w_\lambda^0} T_d c_{A|B} T_{\sigma_0} + \sum_{\substack{u \in \mathfrak{S}_\lambda, \sigma \in \mathcal{D}_{\nu_{A|B}} \cap \mathfrak{S}_\mu \\ u < w_\lambda^0 \text{ or } \sigma < \sigma_0}} T_u T_d c_{A|B} T_\sigma \\ &= T_{w_\lambda^0} T_d c_{A|B} T_{\sigma_0} + \mathfrak{l}_{A|B}, \end{aligned} \quad (5.1.5)$$

where  $\mathfrak{l}_{A|B}$  is an  $R$ -linear combination of the elements  $T_{ud}c_{A|B}T_\sigma$  with  $u \in \mathfrak{S}_\lambda, \sigma \in \mathcal{D}_{\nu(d)} \cap \mathfrak{S}_\mu$  and  $\ell(ud\sigma) < \ell(w_\lambda^0 d\sigma_0) = \ell(w_\lambda^0) + \ell(d) + \ell(\sigma_0)$ . This together with (5.1.4) gives rise to

$$\sum_{(A|B) \in M_n(\mathbb{N}|\mathbb{Z}_2)_{\lambda,\mu}^d} f_{A|B} T_{w_\lambda^0 d} c_{A|B} T_{\sigma_0} + \sum_{(A|B) \in M_n(\mathbb{N}|\mathbb{Z}_2)_{\lambda,\mu}^d} f_{A|B} \mathfrak{l}_{A|B} = 0. \quad (5.1.6)$$

By Lemma 3.7, one can deduce that

$$\sum_{(A|B) \in M_n(\mathbb{N}|\mathbb{Z}_2)_{\lambda,\mu}^d} f_{A|B} T_{w_\lambda^0 d} c_{A|B} T_{\sigma_0} = 0. \quad (5.1.7)$$

Recall the bijection  $\mathfrak{j}$  in (5.0.1). Let  $M^d = (m_{ij}^d) \in M_n(\mathbb{N})$  be the unique matrix such that  $\mathfrak{j}(M^d) = (\lambda, d, \mu)$ . Then  $A + B = M^d$  for all  $(A|B) \in M_n(\mathbb{N}|\mathbb{Z}_2)_{\lambda,\mu}^d$  and we have a bijection between the sets  $M_n(\mathbb{N}|\mathbb{Z}_2)_{\lambda,\mu}^d$  and  $\{B \in M_n(\mathbb{Z}_2) \mid \alpha_B \leq \nu(d)\}$  where  $\nu(d) = (m_{11}^d, \dots, m_{n1}^d, \dots, m_{1n}^d, \dots, m_{nn}^d)$  and  $\alpha_B$  as defined in (5.0.3). Now, it is easy to see that the set  $\{c_{A|B} = c_{\nu(d)}^{\alpha_B} \mid B \in M_n(\mathbb{Z}_2), \alpha_B \leq \nu(d)\}$  is linearly independent in  $\mathcal{C}_r$ . Therefore, by (5.1.7) and Lemma 3.7 again, we have  $f_{A|B} = 0$  for all  $(A|B) \in M_n(\mathbb{N}|\mathbb{Z}_2)_{\lambda,\mu}^d$ .  $\square$

**Proposition 5.2.** *Suppose  $\lambda, \mu \in \Lambda(n, r)$ . Then the intersection  $x_\lambda \mathcal{H}_{r,R}^c \cap \mathcal{H}_{r,R}^c x_\mu$  is a free  $R$ -module with basis*

$$\{T_{A|B} \mid (A|B) \in M_n(\mathbb{N}|\mathbb{Z}_2)_{\lambda,\mu}\}.$$

*Proof.* By Lemma 5.1, it suffices to show that the set spans  $x_\lambda \mathcal{H}_{r,R}^c \cap \mathcal{H}_{r,R}^c x_\mu$ . Take an arbitrary  $z \in x_\lambda \mathcal{H}_{r,R}^c \cap \mathcal{H}_{r,R}^c x_\mu$ . By Corollary 3.8, we may write

$$z = \sum_{(d,\sigma,\alpha) \in X_{\lambda,\mu}} a_{d,\sigma,\alpha} x_\lambda T_d c^\alpha T_\sigma \quad \text{for some } a_{d,\sigma,\alpha} \in R.$$

This together with Corollary 3.5 gives rise to

$$z = \sum_d \left( \sum_{u' \in \mathcal{D}_{\nu(d-1)}^{-1} \cap \mathfrak{S}_\lambda} T_{u'} \right) T_d \sum_{\sigma,\alpha} a_{d,\sigma,\alpha} x_{\nu(d)} c^\alpha T_\sigma.$$

For each  $d \in \mathcal{D}_{\lambda,\mu}$ , write

$$z_d = \sum_{\sigma \in \mathcal{D}_{\nu(d)} \cap \mathfrak{S}_\mu, \alpha \in \mathbb{Z}_2^r} a_{d,\sigma,\alpha} x_{\nu(d)} c^\alpha T_\sigma \in \mathcal{H}_{\mu,R}^c. \quad (5.2.1)$$

Then

$$z = \sum_{d \in \mathcal{D}_{\lambda,\mu}} \sum_{u' \in \mathcal{D}_{\nu(d-1)}^{-1} \cap \mathfrak{S}_\lambda} T_{u'} T_d z_d. \quad (5.2.2)$$

Since  $z \in \mathcal{H}_{r,R}^c x_\mu$ , by Lemma 3.1 we have  $z T_{s_k} = qz$  for  $s_k \in \mathfrak{S}_\mu$  and hence,

$$\sum_{d \in \mathcal{D}_{\lambda,\mu}} \sum_{u' \in \mathcal{D}_{\nu(d-1)}^{-1} \cap \mathfrak{S}_\lambda} T_{u'} T_d (z_d T_{s_k} - qz_d) = 0 \quad (5.2.3)$$

Since  $z_d, z_d T_{s_k} \in \mathcal{H}_{\mu,R}^c$  by (5.2.1) for  $s_k \in \mathfrak{S}_\mu$ , one can deduce that

$$T_{u'} T_d (z_d T_{s_k} - qz_d) \in T_{u'} T_d \mathcal{H}_{\mu,R}^c. \quad (5.2.4)$$

By the second direct sum decomposition in Corollary 3.8, we obtain  $T_{u'} T_d (z_d T_{s_k} - qz_d) = 0$ . Writing  $z_d T_{s_k} - qz_d$  as a linear combination of  $T_w c^\alpha$  with  $w \in \mathfrak{S}_\mu, \alpha \in \mathbb{Z}_2^r$ , linear

independence of  $T_{u'}T_dT_w c^\alpha$  (Lemma 2.3) implies that  $z_d T_{s_k} - q z_d = 0$ , for  $s_k \in \mathfrak{S}_\mu$  and  $d \in \mathcal{D}_{\lambda,\mu}$ . By Lemma 3.1 again, we have

$$z_d \in \mathcal{H}_{r,R}^c x_\mu \cap \mathcal{H}_{\mu,R}^c = \mathcal{H}_{\mu,R}^c x_\mu.$$

Thus, we can write  $z_d$  for each  $d \in \mathcal{D}_{\lambda,\mu}$  as

$$z_d = \sum_{\beta \in \mathbb{Z}_2^r} f_\beta^d c^\beta x_\mu = \sum_{\beta \in \mathbb{Z}_2^r} f_\beta^d c^\beta x_{\nu(d)} \sum_{\sigma \in \mathcal{D}_{\nu(d)} \cap \mathfrak{S}_\mu} T_\sigma$$

for some  $f_\beta^d \in R$  by (5.1.2). Hence, by (5.2.1), we obtain

$$\sum_{\sigma \in \mathcal{D}_{\nu(d)} \cap \mathfrak{S}_\mu, \alpha \in \mathbb{Z}_2^r} a_{d,\sigma,\alpha} x_{\nu(d)} c^\alpha T_\sigma = z_d = \sum_{\beta \in \mathbb{Z}_2^r} f_\beta^d c^\beta x_{\nu(d)} \sum_{\sigma \in \mathcal{D}_{\nu(d)} \cap \mathfrak{S}_\mu} T_\sigma.$$

This means

$$\sum_{\sigma \in \mathcal{D}_{\nu(d)} \cap \mathfrak{S}_\mu} \left( \sum_{\alpha \in \mathbb{Z}_2^r} a_{d,\sigma,\alpha} x_{\nu(d)} c^\alpha - \sum_{\beta \in \mathbb{Z}_2^r} f_\beta^d c^\beta x_{\nu(d)} \right) T_\sigma = 0.$$

But the left hand side belongs to  $\mathcal{H}_{\mu,R}^c = \bigoplus_{\sigma \in \mathcal{D}_{\nu(d)} \cap \mathfrak{S}_\mu} \mathcal{H}_{\nu(d),R}^c T_\sigma$ , forcing every summand is 0. Consequently,

$$\sum_{\alpha \in \mathbb{Z}_2^r} a_{d,\sigma,\alpha} x_{\nu(d)} c^\alpha - \sum_{\beta \in \mathbb{Z}_2^r} f_\beta^d c^\beta x_{\nu(d)} = 0 \quad (5.2.5)$$

for each  $\sigma \in \mathcal{D}_{\nu(d)} \cap \mathfrak{S}_\mu$ . Therefore,

$$\sum_{\alpha \in \mathbb{Z}_2^r} a_{d,\sigma,\alpha} x_{\nu(d)} c^\alpha = \sum_{\beta \in \mathbb{Z}_2^r} f_\beta^d c^\beta x_{\nu(d)} \in x_{\nu(d)} \mathcal{H}_{\nu(d),R}^c \cap \mathcal{H}_{\nu(d),R}^c x_{\nu(d)} \quad (5.2.6)$$

for each  $d \in \mathcal{D}_{\lambda,\mu}$  and  $\sigma \in \mathcal{D}_{\nu(d)} \cap \mathfrak{S}_\mu$ .

On the other hand, since  $d \in \mathcal{D}_{\lambda,\mu}$ , there exists a unique  $M^d = (m_{ij}^d) \in M_n(\mathbb{N})$  such that  $j(M^d) = (\lambda, d, \mu)$  with  $\nu(d) = (m_{11}^d, \dots, m_{n1}^d, \dots, m_{1n}^d, \dots, m_{nn}^d)$ . (Recall  $\nu(d)$  is defined by  $\mathfrak{S}_{\nu(d)} = d^{-1} \mathfrak{S}_\lambda d \cap \mathfrak{S}_\mu$ .) Then, by (5.2.6) and Corollary 4.3,

$$\sum_{\alpha \in \mathbb{Z}_2^r} a_{d,\sigma,\alpha} x_{\nu(d)} c^\alpha = \sum_{B \in M_n(\mathbb{Z}_2), B \leq M^d} g_B^d x_{\nu(d)} c_{\nu(d)}^{\alpha_B} \quad (5.2.7)$$

for some  $g_B^d \in R$ , where  $B = (b_{ij}) \in M_n(\mathbb{Z}_2) \leq M^d$  means  $b_{ij} \leq m_{ij}^d$  for  $1 \leq i, j \leq n$  and  $\alpha_B = (b_{11}, \dots, b_{n1}, \dots, b_{1n}, \dots, b_{nn}) \in \mathbb{Z}_2^{n^2}$ .

For each  $B \in M_n(\mathbb{Z}_2)$  with  $B \leq M^d$ , we let  $A(d, B) = M^d - B \in M_n(\mathbb{Z}_2)$  and then  $d = d_{A(d,B)+B}$  and  $(A(d, B)|B) \in M_n(\mathbb{N}|\mathbb{Z}_2)_{\lambda,\mu}^d$ . Then by (5.0.3) and (5.2.7) we obtain

$$\sum_{\alpha \in \mathbb{Z}_2^r} a_{d,\sigma,\alpha} x_{\nu(d)} c^\alpha = \sum_{B \in M_n(\mathbb{Z}_2), B \leq M^d} g_B^d x_{\nu(d)} c_{A(d,B)|B}$$

and hence, by (5.2.1),

$$z_d = \sum_{\sigma \in \mathcal{D}_{\nu(d)} \cap \mathfrak{S}_\mu} \sum_{B \in M_n(\mathbb{Z}_2), B \leq M^d} g_B^d x_{\nu(d)} c_{A(d,B)|B} T_\sigma.$$

Therefore, by (5.2.2), one can deduce that

$$\begin{aligned}
z &= \sum_{d \in \mathcal{D}_{\lambda, \mu}} \sum_{u' \in \mathcal{D}_{\nu(d-1)}^{-1} \cap \mathfrak{S}_{\lambda}} T_{u'} T_d \sum_{\sigma \in \mathcal{D}_{\nu(d)} \cap \mathfrak{S}_{\mu}} \sum_{B \in M_n(\mathbb{Z}_2), B \leq M^d} g_B^d x_{\nu(d)} c_{A(d, B)|B} T_{\sigma} \\
&= \sum_{d \in \mathcal{D}_{\lambda, \mu}} \sum_{B \in M_n(\mathbb{Z}_2), B \leq M^d} g_B^d \sum_{u' \in \mathcal{D}_{\nu(d-1)}^{-1} \cap \mathfrak{S}_{\lambda}} T_{u'} T_d x_{\nu(d)} c_{A(d, B)|B} \sum_{\sigma \in \mathcal{D}_{\nu(d)} \cap \mathfrak{S}_{\mu}} T_{\sigma} \\
&= \sum_{d \in \mathcal{D}_{\lambda, \mu}} \sum_{B \in M_n(\mathbb{Z}_2), B \leq M^d} g_B^d x_{\lambda} T_d c_{A(d, B)|B} \sum_{\sigma \in \mathcal{D}_{\nu(d)} \cap \mathfrak{S}_{\mu}} T_{\sigma} \\
&= \sum_{d \in \mathcal{D}_{\lambda, \mu}} \sum_{B \in M_n(\mathbb{Z}_2), B \leq M^d} g_B^d T_{A(d, B)|B}
\end{aligned}$$

since  $d = d_{A(d, B)+B}$ . This proves the proposition.  $\square$

For  $(A|B) \in M_n(\mathbb{N}|\mathbb{Z}_2)_r$ , define  $\phi_{(A|B)} \in \mathcal{Q}_q(n, r; R) = \text{End}_{\mathcal{H}_{r, R}^c}(\oplus_{\mu \in \Lambda(n, r)} x_{\mu} \mathcal{H}_{r, R}^c)$  via

$$\phi_{(A|B)}(x_{\mu} h) = \delta_{\mu, \text{co}(A+B)} T_{A|B} h \quad (5.2.8)$$

for  $\mu \in \Lambda(n, r)$  and  $h \in \mathcal{H}_{r, R}^c$ .

**Theorem 5.3.** *Let  $R$  be a commutative ring of characteristic not equal to 2. Then the algebra  $\mathcal{Q}_q(n, r; R)$  is a free  $R$ -module with a basis given by the set*

$$\{\phi_{(A|B)} \mid (A|B) \in M_n(\mathbb{N}|\mathbb{Z}_2)_r\}.$$

*Proof.* By (2.4.3), one can deduce that

$$\mathcal{Q}_q(n, r; R) = \bigoplus_{\lambda, \mu \in \Lambda(n, r)} \text{Hom}_{\mathcal{H}_{r, R}^c}(x_{\mu} \mathcal{H}_{r, R}^c, x_{\lambda} \mathcal{H}_{r, R}^c).$$

Then, by Lemma 3.2, Proposition 5.2, the set

$$\{\phi_{(A|B)} \mid \text{ro}(A+B) = \lambda, \text{co}(A+B) = \mu\}$$

is an  $R$ -basis for  $\text{Hom}_{\mathcal{H}_{r, R}^c}(x_{\mu} \mathcal{H}_{r, R}^c, x_{\lambda} \mathcal{H}_{r, R}^c)$  for each pair  $\lambda, \mu \in \Lambda(n, r)$ . Therefore, the set

$$\{\phi_{(A|B)} \mid (A|B) \in M_n(\mathbb{N}|\mathbb{Z}_2)_r\}$$

forms an  $R$ -basis for  $\mathcal{Q}_q(n, r; R)$ .  $\square$

By Theorem 5.3, we have the following base change property for  $\mathcal{Q}_q(n, r; R)$ .

**Corollary 5.4.** *Maintain the assumption on  $R$  as above. Suppose that  $R$  is an  $\mathcal{A}$ -algebra via  $\mathbf{q} \mapsto q$ . Then*

$$\mathcal{Q}_q(n, r; R) \cong \mathcal{Q}_{\mathbf{q}}(n, r)_R := \mathcal{Q}_{\mathbf{q}}(n, r) \otimes_{\mathcal{A}} R.$$

## 6. IDENTIFICATION WITH THE QUOTIENTS OF THE QUANTUM QUEER SUPERGROUP

In this section, we shall show that the  $\mathbf{q}$ - $q$ -Schur superalgebra  $\mathcal{Q}_{\mathbf{v}}(n, r)$  coincides with the quantum queer Schur superalgebra constructed in [12]. In particular, they are homomorphic images of the quantum queer supergroup  $U_{\mathbf{v}}(\mathbf{q}_n)$ .

Let  $q = v^2$  for some  $v \in R$ . Let  $V(n|n)_R = V_0 \oplus V_1$  be a free  $R$ -supermodule with basis  $e_1, \dots, e_n$  for  $V_0$  and basis  $e_{-1}, \dots, e_{-n}$  for  $V_1$ . With the fixed ordered basis  $\{e_1, \dots, e_n, e_{-1}, \dots, e_{-n}\}$ , we often identify the  $R$ -algebra  $\text{End}_R(V(n|n)_R)$  of all  $R$ -linear maps on  $V(n|n)_R$  with the  $2n \times 2n$  matrix algebra  $M_{2n}(R)$  over  $R$  and, hence,  $\text{End}_R(V(n|n)_R^{\otimes r})$  with  $M_{2n}(R)^{\otimes r}$ .

Let

$$I(n|n) = \{1, 2, \dots, n, -1, -2, \dots, -n\},$$

$$I(n|n, r) = \{\underline{i} = (i_1, \dots, i_r) \mid i_k \in I(n|n), 1 \leq k \leq r\}.$$

Then,  $\mathfrak{S}_r$  acts on  $I(n|n, r)$  by place permutation

$$\underline{i}s_k = (i_1, \dots, i_{k-1}, i_{k+1}, i_k, i_{k+2}, \dots, i_r).$$

For  $\underline{i} = (i_1, \dots, i_r) \in I(n|n, r)$ , set  $e_{\underline{i}} = e_{i_1} \otimes e_{i_2} \otimes \dots \otimes e_{i_r}$ , then the set  $\{e_{\underline{i}} \mid \underline{i} \in I(n|n, r)\}$  forms a basis for  $V(n|n)_R^{\otimes r}$ .

Denote by  $E_{i,j}$  for  $i, j \in I(n|n)$  the standard elementary matrix with the  $(i, j)$ th entry being 1 and 0 elsewhere. Then  $\{E_{i,j} \mid i, j \in I(n|n)\}$  can be viewed as the standard basis for  $\text{End}_R(V(n|n)_R)$ , that is,  $E_{i,j}(e_k) = \delta_{j,k}e_i$ . Following [20], we set

$$\begin{aligned} \Omega &= \sum_{1 \leq a \leq n} (E_{-a,a} - E_{a,-a}), \\ T &= \sum_{i,j \in I(n|n)} (-1)^{\widehat{j}} E_{i,j} \otimes E_{j,i}, \\ S &= v \sum_{i \leq j \in I(n|n)} S_{i,j} \otimes E_{i,j} \in \text{End}_R(V(n|n)_R^{\otimes 2}), \end{aligned} \tag{6.0.1}$$

where  $\widehat{j} = \widehat{e}_j = 0$  if  $j > 0$  and  $\widehat{j} = \widehat{e}_j = 1$  if  $j < 0$ , and  $S_{i,j}$  for  $i \leq j$  are defined as follows:

$$\begin{aligned} S_{a,a} &= 1 + (v - 1)(E_{a,a} + E_{-a,-a}), \quad 1 \leq a \leq n, \\ S_{-a,-a} &= 1 + (v^{-1} - 1)(E_{a,a} + E_{-a,-a}), \quad 1 \leq a \leq n, \\ S_{b,a} &= (v - v^{-1})(E_{a,b} + E_{-a,-b}), \quad 1 \leq b < a \leq n, \\ S_{-b,-a} &= -(v - v^{-1})(E_{a,b} + E_{-a,-b}), \quad 1 \leq a < b \leq n, \\ S_{-b,a} &= -(v - v^{-1})(E_{-a,b} + E_{a,-b}), \quad 1 \leq a, b \leq n. \end{aligned} \tag{6.0.2}$$

To endomorphisms  $A \in \text{End}_R(V(n|n)_R)$  and  $Z = \sum_t X_t \otimes Y_t \in \text{End}_R(V(n|n)_R)^{\otimes 2} = \text{End}_R(V(n|n)_R^{\otimes 2})$ , we associate the following elements in  $\text{End}_R(V(n|n)_R^{\otimes r})$ :

$$\begin{aligned} A^{(k)} &= \text{id}^{\otimes(k-1)} \otimes A \otimes \text{id}^{\otimes(r-k)}, \quad 1 \leq k \leq r, \\ Z^{(j,k)} &= \sum_t X_t^{(j)} Y_t^{(k)}, \quad 1 \leq j \neq k \leq r. \end{aligned}$$

Let  $\bar{S} = TS$ . It follows from [20, Theorems 5.2-5.3] that there exists a *left*  $\mathcal{H}_{r,R}^c$ -supermodule structure on  $V(n|n)_R^{\otimes r}$  given by

$$\Psi_r : \mathcal{H}_{r,R}^c \longrightarrow \text{End}_R(V(n|n)_R^{\otimes r}), \quad T_i \longmapsto \bar{S}^{(i,i+1)}, c_j \longmapsto \Omega^{(j)} \tag{6.0.3}$$

for all  $1 \leq i \leq r-1, 1 \leq j \leq r$ . Here we remark that the even generators  $T_1, \dots, T_{r-1}$  for  $\mathcal{H}_{r,R}^c$  are related to the even generators  $t_1, \dots, t_{r-1}$  in [20] via  $T_i = vt_i$  for  $1 \leq i \leq r-1$ . This above action has been explicitly worked out in [22, Lemma 3.1] via (6.0.1):

**Lemma 6.1.** *The  $\mathcal{H}_{r,R}^c$ -supermodule structure on  $V(n|n)_R^{\otimes r}$  is given by the following formulas: for  $\underline{i} = (i_1, \dots, i_r) \in I(n|n, r)$  and  $1 \leq k \leq r-1$ ,*

$$c_j \cdot e_{\underline{i}} = (-1)^{\hat{e}_{i_1} + \dots + \hat{e}_{i_j}} e_{i_1} \otimes \dots \otimes e_{i_{j-1}} \otimes e_{-i_j} \otimes e_{i_{j+1}} \otimes \dots \otimes e_{i_r}.$$

$$T_k \cdot e_{\underline{i}} = \begin{cases} v^2 e_{\underline{i}s_k} + (v^2 - 1) e_{\underline{i}_k^-}, & \text{if } i_k = i_{k+1} \geq 1, \\ -e_{\underline{i}s_k}, & \text{if } i_k = i_{k+1} \leq -1, \\ e_{\underline{i}s_k}, & \text{if } i_k = -i_{k+1} \geq 1, \\ v^2 e_{\underline{i}s_k} + (v^2 - 1) e_{\underline{i}}, & \text{if } i_k = -i_{k+1} \leq -1, \\ v e_{\underline{i}s_k} + (v^2 - 1) e_{\underline{i}_k^-} + (v^2 - 1) e_{\underline{i}}, & \text{if } |i_k| < |i_{k+1}| \text{ and } i_{k+1} \geq 1, \\ (-1)^{\hat{i}_k} v e_{\underline{i}s_k}, & \text{if } |i_k| < |i_{k+1}| \text{ and } i_{k+1} \leq -1, \\ v e_{\underline{i}s_k} + (-1)^{\hat{i}_{k+1}} (v^2 - 1) e_{\underline{i}_k^-}, & \text{if } |i_{k+1}| < |i_k| \text{ and } i_k \geq 1, \\ (-1)^{\hat{i}_{k+1}} e_{\underline{i}s_k} + (v^2 - 1) e_{\underline{i}}, & \text{if } |i_{k+1}| < |i_k| \text{ and } i_k \leq -1, \end{cases}$$

where  $\underline{i}_k^- = (i_1, \dots, i_{k-1}, -i_k, -i_{k+1}, i_{k+2}, \dots, i_r)$ .

For each  $\underline{i} \in I(n|n, r)$ , define  $\text{wt}(\underline{i}) = \lambda = (\lambda_1, \dots, \lambda_n) \in \Lambda(n, r)$  by setting

$$\lambda_a = \#\{k \mid a = |i_k|, 1 \leq k \leq r\}, \quad \forall 1 \leq a \leq n.$$

For each  $\lambda \in \Lambda(n, r)$ , define  $\underline{i}_\lambda$  by

$$\underline{i}_\lambda = (\underbrace{-1, \dots, -1}_{\lambda_1}, \underbrace{-2, \dots, -2}_{\lambda_2}, \dots, \underbrace{-n, \dots, -n}_{\lambda_n}).$$

Recall from (2.4.2) the elements  $y_\lambda$ .

**Corollary 6.2.** *The following holds as left  $\mathcal{H}_{r,R}^c$ -supermodules:*

$$V(n|n)_R^{\otimes r} \cong \bigoplus_{\lambda \in \Lambda(n, r)} \mathcal{H}_{r,R}^c y_\lambda.$$

*Proof.* For each  $\lambda \in \Lambda(n, r)$ , let  $(V(n|n)_R^{\otimes r})_\lambda$  be the  $R$ -submodule of  $V(n|n)_R^{\otimes r}$  spanned by the elements  $e_{\underline{i}}$  with  $\underline{i} \in I(n|n, r)$  and  $\text{wt}(\underline{i}) = \lambda$ . Then, by Lemma 6.1,  $(V(n|n)_R^{\otimes r})_\lambda$  is stable under the action of  $\mathcal{H}_{r,R}^c$  and, moreover,  $V(n|n)_R^{\otimes r}$  can be decomposed as

$$V(n|n)_R^{\otimes r} = \bigoplus_{\lambda \in \Lambda(n, r)} (V(n|n)_R^{\otimes r})_\lambda. \quad (6.2.1)$$

Clearly  $e_{\underline{i}_\lambda} \in (V(n|n)_R^{\otimes r})_\lambda$ . For each  $w \in \mathfrak{S}_r$ , we have  $\underline{i}_\lambda w = (i_{w(1)}, \dots, i_{w(r)})$ , where we write  $\underline{i}_\lambda = (i_1, \dots, i_r)$ . Then we can easily deduce that

$$\{e_{\underline{i}} \mid \underline{i} \in I(n|n, r), \text{wt}(\underline{i}) = \lambda, i_k \leq -1, 1 \leq k \leq r\} = \{e_{\underline{i}_\lambda w} \mid w \in \mathfrak{S}_r\} = \{e_{\underline{i}_\lambda d} \mid d \in \mathcal{D}_\lambda\}.$$

Moreover by Lemma 6.1, one can deduce that

$$\mathcal{H}_{r,R} e_{\underline{i}_\lambda} = R\text{-span}\{e_{\underline{i}_\lambda d} \mid d \in \mathcal{D}_\lambda\}.$$

Meanwhile by Frobenius reciprocity there exists an  $\mathcal{H}_{r,R}^c$ -homomorphism

$$\mathcal{H}_{r,R}^c \otimes_{\mathcal{H}_{r,R}} \mathcal{H}_{r,R} e_{\underline{i}_\lambda} \rightarrow \mathcal{H}_{r,R}^c e_{\underline{i}_\lambda} = (V(n|n)_R^{\otimes r})_\lambda, \quad c^\alpha \otimes e_{\underline{i}_\lambda d} \mapsto c^\alpha \cdot e_{\underline{i}_\lambda d} \quad (6.2.2)$$

which is bijective due to the fact that  $\{c^\alpha \cdot e_{\underline{i}_\lambda d} \mid d \in \mathcal{D}_\lambda, \alpha \in \mathbb{Z}_2^r\}$  is a basis for  $(V(n|n)_R^{\otimes r})_\lambda$ . Observe that the tensor product  $V(0|n)_R^{\otimes r}$  of odd  $R$ -submodule  $V(0|n)_R$  of  $V(n|n)_R$  admits a basis  $\{e_{\underline{i}} \mid \underline{i} \in I(n|n, r), i_k \leq -1, 1 \leq k \leq r\}$ , which can be identified with

$\bigoplus_{\lambda \in \Lambda(n,r)} \mathcal{H}_{r,R} e_{\underline{i}_\lambda}$ . Then, by Lemma 6.1 and [8, Lemma 2.1], there is an  $\mathcal{H}_{r,R}$ -module isomorphism

$$\mathcal{H}_{r,R} e_{\underline{i}_\lambda} \cong \mathcal{H}_{r,R} y_\lambda. \quad (6.2.3)$$

Taking a direct sum over  $\Lambda(n,r)$  and induction to  $\mathcal{H}_{r,R}^c$  and noting (6.2.1) and (6.2.2) gives the required isomorphism.  $\square$

Recall the anti-involution  $\tau$  defined in Lemma 2.2(3) and the twist functor in Remark 2.5. In particular,  $(V(n|n)_R^{\otimes r})^\tau$  affords a right  $\mathcal{H}_{r,R}^c$ -supermodule. By (2.5.1), we immediately have the following.

**Corollary 6.3.** *There is an isomorphism of right  $\mathcal{H}_{r,R}^c$ -supermodules:*

$$(V(n|n)_R^{\otimes r})^\tau \cong \bigoplus_{\lambda \in \Lambda(n,r)} x_\lambda \mathcal{H}_{r,R}^c,$$

which induces an algebra isomorphism

$$\text{End}_{\mathcal{H}_{r,R}^c}(V(n|n)_R^{\otimes r}) \cong \text{End}_{\mathcal{H}_{r,R}^c}((V(n|n)_R^{\otimes r})^\tau) \cong \text{End}_{\mathcal{H}_{r,R}^c} \left( \bigoplus_{\lambda \in \Lambda(n,r)} x_\lambda \mathcal{H}_{r,R}^c \right) = \mathcal{Q}_{v^2}(n, r; R).$$

Hence, if  $R = \mathcal{Z} = \mathbb{Z}[\mathbf{v}, \mathbf{v}^{-1}]$ , the endomorphism superalgebra  $\text{End}_{\mathcal{H}_{r,R}^c}(V(n|n)_R^{\otimes r})$  is the quantum queer Schur superalgebra  $\mathcal{Q}_{\mathbf{v}}(n, r)$  considered in [12].<sup>1</sup> The isomorphism above shows that our notation is consistent as, by (2.4.4), we used the same notation to denote the right hand side  $\mathcal{Q}_{v^2}(n, r; \mathcal{Z})$ ; see (2.4.4).

Olshanski [20] introduced the quantum deformation  $U_{\mathbf{v}}(\mathfrak{q}_n)$  over  $\mathbb{C}(\mathbf{v})$  of the universal enveloping algebra of the queer Lie superalgebra  $\mathfrak{q}(n)$  and defined a superalgebra homomorphism

$$\Phi_r : U_{\mathbf{v}}(\mathfrak{q}_n) \longrightarrow \text{End}_{\mathbb{C}(\mathbf{v})}(V(n|n)_{\mathbb{C}(\mathbf{v})}^{\otimes r}).$$

We refer the reader to [20] for the details of the definitions of  $U_{\mathbf{v}}(\mathfrak{q}(n))$  and  $\Phi_r$ . The following result is known as the double centraliser property and forms the first part of the quantum analog of Schur-Weyl–Sergeev duality for  $U_{\mathbf{v}}(\mathfrak{q}_n)$  and  $\mathcal{H}_{r,\mathbb{C}(\mathbf{v})}^c$ . Recall also the algebra homomorphism  $\Psi_r$  defined in (6.0.3).

**Proposition 6.4** ([20, Theorem 5.3]). *The algebras  $\Phi_r(U_{\mathbf{v}}(\mathfrak{q}_n))$  and  $\Psi_r(\mathcal{H}_{r,\mathbb{C}(\mathbf{v})}^c)$  form mutual centralisers in  $\text{End}_{\mathbb{C}(\mathbf{v})}(V(n|n)_{\mathbb{C}(\mathbf{v})}^{\otimes r})$ . In particular,  $\Phi_r$  induces an algebra epimorphism*

$$\Phi_r : U_{\mathbf{v}}(\mathfrak{q}_n) \longrightarrow \text{End}_{\mathcal{H}_{r,\mathbb{C}(\mathbf{v})}^c}(V(n|n)_{\mathbb{C}(\mathbf{v})}^{\otimes r}) = \mathcal{Q}_{\mathbf{v}}(n, r)_{\mathbb{C}(\mathbf{v})}.$$

Thus, by [12, Theorem 9.2], the  $\mathbb{Q}$ - $q$ -Schur superalgebra  $\mathcal{Q}_{\mathbf{v}}(n, r)_{\mathbb{C}(\mathbf{v})}$  has a presentation with even generators  $K_i^{\pm 1}, E_j, F_j$  and odd generators  $K_{\bar{i}}, E_{\bar{j}}, F_{\bar{j}}$ , for  $1 \leq i \leq n$  and  $1 \leq j \leq n-1$ , subject to the relations (QQ1)–(QQ9) given in [12].

<sup>1</sup>In [12], we used the indeterminate  $q$  instead of  $\mathbf{v}$  and the notation  $\mathcal{Q}_q(n, r)$  denotes the queer  $q$ -Schur superalgebra over the rational function field  $\mathbb{C}(q)$ .

7. IRREDUCIBLE  $\mathcal{Q}_v(n, r)_{\mathbb{K}}$ -SUPERMODULES

In this section, we shall give a classification of irreducible  $\mathcal{Q}_v(n, r)_{\mathbb{K}}$ -supermodules over certain field extension  $\mathbb{K}$  of  $\mathcal{Z} = \mathbb{Z}[\mathbf{v}, \mathbf{v}^{-1}]$ . But we first look at a few general facts for the algebra  $\mathcal{Q}_q(n, r)$  over  $\mathcal{A} = \mathbb{Z}[\mathbf{q}]$ .

Let  $\omega = (1^r)$ . In the case  $n \geq r$ , we can view  $\omega$  as an element in  $\Lambda(n, r)$  and define

$$e_\omega = \phi_{(A_\omega | B_\omega)} \in \mathcal{Q}_q(n, r),$$

where  $A_\omega = \text{diag}(\underbrace{1, \dots, 1}_r, 0, \dots, 0)$  and  $B_\omega = 0$ . Then, by (5.0.4) and (5.2.8),

$$e_\omega(x_\lambda h) = \delta_{\omega, \lambda} x_\lambda h, \quad e_\omega^2 = e_\omega \quad (7.0.1)$$

for  $\lambda \in \Lambda(n, r)$ ,  $h \in \mathcal{H}_r^c$ .

Recall from [7, Theorem 2.24] that the  $q$ -Schur algebra  $\mathcal{S}_q(n, r)$  is defined as

$$\mathcal{S}_q(n, r) := \text{End}_{\mathcal{H}_r} \left( \bigoplus_{\lambda \in \Lambda(n, r)} x_\lambda \mathcal{H}_r \right). \quad (7.0.2)$$

Let

$$\mathcal{T}(n, r) = \bigoplus_{\lambda \in \Lambda(n, r)} x_\lambda \mathcal{H}_r \quad \text{and} \quad \mathcal{T}^c(n, r) = \bigoplus_{\lambda \in \Lambda(n, r)} x_\lambda \mathcal{H}_r^c.$$

Since  $\mathcal{T}^c(n, r) = \mathcal{T}(n, r) \otimes_{\mathcal{H}_r} \mathcal{H}_r^c$  (see (2.4.5)), there is a natural algebra embedding

$$\epsilon : \mathcal{S}_q(n, r) \longrightarrow \mathcal{Q}_q(n, r), f \longmapsto f \otimes 1.$$

Equivalently, the  $q$ -Schur algebra  $\mathcal{S}_q(n, r)$  can be identified as a subalgebra of  $\mathcal{Q}_q(n, r)$  via the following way

$$\mathcal{S}_q(n, r) = \{ \phi \in \mathcal{Q}_q(n, r) \mid \phi(x_\lambda) \in \mathcal{H}_r, \forall \lambda \in \Lambda(n, r) \}. \quad (7.0.3)$$

On the other hand, if  $n \geq r$ , the evaluation map gives algebra isomorphisms

$$e_\omega \mathcal{Q}_q(n, r) e_\omega \cong \mathcal{H}_r^c \quad \text{and} \quad e_\omega \mathcal{S}_q(n, r) e_\omega \cong \mathcal{H}_r.$$

We will identify them in sequel. In particular,  $\mathcal{Q}_q(n, r) e_\omega$  is a  $\mathcal{Q}_q(n, r)$ - $\mathcal{H}_r^c$ -bisupermodule.

**Lemma 7.1.** *If  $n \geq r$ , then there is a  $\mathcal{Q}_q(n, r)$ - $\mathcal{H}_r^c$ -bisupermodule isomorphism*

$$\mathcal{Q}_q(n, r) e_\omega \cong \mathcal{T}^c(n, r).$$

*Moreover, restriction gives an  $\mathcal{S}_q(n, r)$ - $\mathcal{H}_r^c$ -bisupermodule isomorphism*

$$\mathcal{Q}_q(n, r) e_\omega \cong \mathcal{S}_q(n, r) e_\omega \otimes_{\mathcal{H}_r} \mathcal{H}_r^c.$$

*Proof.* By (7.0.1) and (7.0.3),  $e_\omega \in \mathcal{S}_q(n, r)$  and  $x_\omega = 1$ . Hence, the evaluation map gives an  $\mathcal{S}_q(n, r)$ - $\mathcal{H}_r$ -bimodule isomorphism

$$\mathcal{S}_q(n, r) e_\omega = \text{Hom}_{\mathcal{H}_r}(x_\omega \mathcal{H}_r, \mathcal{T}(n, r)) \cong \mathcal{T}(n, r),$$

and a  $\mathcal{Q}_q(n, r)$ - $\mathcal{H}_r^c$ -bisupermodule isomorphism

$$\mathcal{Q}_q(n, r) e_\omega = \text{Hom}_{\mathcal{H}_r^c}(x_\omega \mathcal{H}_r^c, \mathcal{T}^c(n, r)) \cong \mathcal{T}^c(n, r).$$

Thus, the restriction gives an  $\mathcal{S}_q(n, r)$ - $\mathcal{H}_r^c$ -bisupermodule isomorphism

$$\mathcal{Q}_q(n, r) e_\omega \cong \mathcal{T}(n, r) \otimes_{\mathcal{H}_r} \mathcal{H}_r^c \cong \mathcal{S}_q(n, r) e_\omega \otimes_{\mathcal{H}_r} \mathcal{H}_r^c.$$

This proves the lemma.  $\square$

Assume that  $\mathbb{F}$  is a field which is a  $\mathcal{Z}$ -algebra such that the image of  $v$  is not a root of unity. Thus, both  $\mathcal{H}_{r,\mathbb{F}}$  and  $\mathcal{S}_q(n, r)_{\mathbb{F}} = \mathcal{S}_q(n, r) \otimes_{\mathcal{Z}} \mathbb{F}$  are semisimple  $\mathbb{F}$ -algebras. The following result generalizes the category equivalence between  $\mathcal{S}_q(n, r)_{\mathbb{F}}\text{-mod}$  and  $\mathcal{H}_{r,\mathbb{F}}\text{-mod}$  in the case  $n \geq r$  and the proof is somewhat standard, see, e.g., [4, Theorem 4.1.3].

**Proposition 7.2.** *Assume  $n \geq r$ . The superalgebras  $\mathcal{Q}_q(n, r)_{\mathbb{F}}$  and  $\mathcal{H}_{r,\mathbb{F}}^c$  are Morita equivalent.*

*Proof.* We consider the two functors generally defined in [14, §6.2]. Firstly, the  $\mathcal{Q}_q(n, r)_{\mathbb{F}}\text{-}\mathcal{H}_{r,\mathbb{F}}^c$ -bisupermodule  $\mathcal{T}^c(n, r)_{\mathbb{F}}$  induces a functor

$$\mathcal{F} : \mathcal{H}_{r,\mathbb{F}}^c\text{-smod} \longrightarrow \mathcal{Q}_q(n, r)_{\mathbb{F}}\text{-smod}, \quad L \longmapsto \mathcal{T}^c(n, r)_{\mathbb{F}} \otimes_{\mathcal{H}_{r,\mathbb{F}}^c} L. \quad (7.2.1)$$

Secondly, there exists the functor

$$\mathcal{G} : \mathcal{Q}_q(n, r)_{\mathbb{F}}\text{-smod} \rightarrow \mathcal{H}_{r,\mathbb{F}}^c\text{-smod}, \quad M \mapsto e_{\omega}M.$$

It is clear (cf. [14, (6.2d)]) that  $\mathcal{G} \circ \mathcal{F} \cong \text{id}_{\mathcal{H}_{r,\mathbb{F}}^c\text{-mod}}$ . So it suffices to prove  $\mathcal{F} \circ \mathcal{G} \cong \text{id}_{\mathcal{Q}_q(n, r)_{\mathbb{F}}\text{-mod}}$ .

It is known from the case for the  $q$ -Schur algebra that, for any  $\mathcal{S}_q(n, r)_{\mathbb{F}}$ -module  $M$ , there is a left  $\mathcal{S}_q(n, r)_{\mathbb{F}}$ -module isomorphism

$$\psi : \mathcal{S}_q(n, r)_{\mathbb{F}} e_{\omega} \otimes_{\mathcal{H}_{r,\mathbb{F}}} e_{\omega} M \cong M$$

defined by  $\psi(x \otimes m) = xm$  for  $x \in \mathcal{S}_q(n, r)_{\mathbb{F}} e_{\omega}$  and  $m \in e_{\omega} M$ . This together with Lemma 7.1 gives rise to a left  $\mathcal{S}_q(n, r)_{\mathbb{F}}$ -module isomorphism

$$\Psi : \mathcal{Q}_q(n, r)_{\mathbb{F}} e_{\omega} \otimes_{\mathcal{H}_{r,\mathbb{F}}^c} e_{\omega} M = \mathcal{S}_q(n, r)_{\mathbb{F}} e_{\omega} \otimes_{\mathcal{H}_{r,\mathbb{F}}} e_{\omega} M \cong M$$

We now claim that  $\Psi$  is a  $\mathcal{Q}_q(n, r)_{\mathbb{F}}$ -supermodule isomorphism. Indeed, considering the basis  $\{\phi_{(A|B)} \mid (A|B) \in M_n(\mathbb{N}|\mathbb{Z}_2)\}$  for  $\mathcal{Q}_q(n, r)_{\mathbb{F}}$ , we have

$$\mathcal{Q}_q(n, r)_{\mathbb{F}} e_{\omega} = \text{Hom}_{\mathcal{H}_{r,\mathbb{F}}^c}(x_{\omega} \mathcal{H}_{r,\mathbb{F}}^c, \bigoplus_{\lambda \in \Lambda(n, r)} x_{\lambda} \mathcal{H}_{r,\mathbb{F}}^c) = \bigoplus_{\lambda \in \Lambda(n, r)} \phi_{(A(\omega, \lambda)|0)} \mathcal{H}_{r,\mathbb{F}}^c,$$

where  $A(\omega, \lambda) = (a_{ij}^{\omega, \lambda})$  with  $a_{ij}^{\omega, \lambda} = 1$  if  $\lambda_1 + \dots + \lambda_{i-1} + 1 \leq j \leq \lambda_1 + \dots + \lambda_{i-1} + \lambda_i$  and  $a_{ij}^{\omega, \lambda} = 0$  otherwise. Clearly  $\phi_{(A(\omega, \lambda)|0)}(x_{\omega}) = x_{\lambda} \in \mathcal{H}_{r,\mathbb{F}}$  for  $\lambda \in \Lambda(n, r)$ . Therefore  $\phi_{(A(\omega, \lambda)|0)} \in \mathcal{S}_q(n, r)_{\mathbb{F}} e_{\omega}$  by (7.0.3). This implies that

$$\Psi(\phi_{(A(\omega, \lambda)|0)} h \otimes m) = \psi(\phi_{(A(\omega, \lambda)|0)} \otimes hm) = \phi_{(A(\omega, \lambda)|0)} hm$$

for any  $h \in \mathcal{H}_{r,\mathbb{F}}^c, m \in e_{\omega} M$ . Hence  $\Psi(x \otimes m) = xm$  for  $x \in \mathcal{Q}_q(n, r)_{\mathbb{F}} e_{\omega}, m \in e_{\omega} M$ . This means  $\Psi$  is a  $\mathcal{Q}_q(n, r)_{\mathbb{F}}$ -supermodule homomorphism. Then the proposition is proved.  $\square$

We are now ready to look at the classification of irreducible representations of the superalgebra  $\mathcal{Q}_v(n, r)_{\mathbb{K}}$ , where

$$\mathbb{K} := \mathbb{C}(\mathbf{v}) \left( \sqrt{[2]}, \sqrt{[3]}, \dots, \sqrt{[r]} \right)$$

is the field extension<sup>2</sup> of  $\mathbb{C}(\mathbf{v})$  with  $[k] = \frac{q^k - q^{-k}}{q - q^{-1}}$  for  $k \in \mathbb{N}$  ( $q = v^2$ ). It is known from [16, Proposition 2.2] that  $\mathcal{H}_{r, \mathbb{C}(\mathbf{v})}^c$  is a semisimple superalgebra. Thus,  $\mathcal{Q}_v(n, r)_{\mathbb{C}(\mathbf{v})}$  is semisimple. The following result will imply that  $\mathbb{K}$  is a splitting field of  $\mathcal{Q}_v(n, r)_{\mathbb{C}(\mathbf{v})}$ .

<sup>2</sup>The field  $\mathbb{K}$  is denoted by  $\mathbb{F}$  in [16, §4].

A partition  $\xi$  of  $r$  is said to be *strict* if its non-zero parts are distinct. Denote by  $\mathcal{SP}(r)$  the set of strict partitions  $\xi$  of  $r$ . For  $\xi \in \mathcal{SP}(r)$ , denote by  $l(\xi)$  the length of  $\xi$  and let

$$\delta(\xi) = \begin{cases} 0, & \text{if } l(\xi) \text{ is even,} \\ 1, & \text{if } l(\xi) \text{ is odd.} \end{cases}$$

**Proposition 7.3** ([16, Corollary 6.8]). *For each  $\xi \in \mathcal{SP}(r)$ , there exists a left simple  $\mathcal{H}_{r,\mathbb{K}}^c$ -supermodule  $U^\xi$  such that  $\{U^\xi \mid \xi \in \mathcal{SP}(r)\}$  forms a complete set of nonisomorphic irreducible left  $\mathcal{H}_{r,\mathbb{K}}^c$ -supermodules. Moreover, every  $U^\xi$  is split irreducible<sup>3</sup> and  $U^\xi$  is of type  $M$  (resp.,  $Q$ ) if  $\delta(\xi) = 0$  (resp.,  $\delta(\xi) = 1$ ).*

If we take  $\mathbb{F} = \mathbb{K}$  in (7.2.1), then Propositions 7.2 and 7.3 imply the following immediately.

**Corollary 7.4.** *If  $n \geq r$ , then the set  $\{\mathcal{T}^c(n, r)_{\mathbb{K}} \otimes_{\mathcal{H}_{r,\mathbb{K}}^c} U^\xi \mid \xi \in \mathcal{SP}(r)\}$  is a complete set of nonisomorphic irreducible  $\mathcal{Q}_v(n, r)_{\mathbb{K}}$ -supermodules.*

For positive integers  $n, r$ , let

$$\mathcal{SP}(n, r) = \{\xi \in \mathcal{SP}(r) \mid \mathcal{T}^c(n, r)_{\mathbb{K}} \otimes_{\mathcal{H}_{r,\mathbb{K}}^c} U^\xi \neq 0\}.$$

Clearly, we have from the above  $\mathcal{SP}(n, r) = \mathcal{SP}(r)$  for all  $n \geq r$ .

Recall from the last section, there is a commuting action of  $U_v(\mathfrak{q}_n)_{\mathbb{K}}$  and  $\mathcal{H}_{r,\mathbb{K}}^c$  on  $V(n|n)_{\mathbb{K}}^{\otimes r}$ . For  $\xi \in \mathcal{SP}(r)$ , let

$$V(\xi) = \text{Hom}_{\mathcal{H}_{r,\mathbb{K}}^c}(U^\xi, V(n|n)_{\mathbb{K}}^{\otimes r}).$$

This is a  $U_v(\mathfrak{q}_n)_{\mathbb{K}}$ -module. If  $V(\xi) \neq 0$ , then  $U^\xi$  is a direct summand of  $V(n|n)_{\mathbb{K}}^{\otimes r}$ . Hence it is a simple  $\mathcal{Q}_v(n, r)_{\mathbb{K}}$ -module and, hence, a simple  $U_v(\mathfrak{q}_n)_{\mathbb{K}}$ -module.

**Lemma 7.5.** *Let  $\xi \in \mathcal{SP}(r)$ . If  $n < r$  and  $l(\xi) > n$ , then  $V(\xi) = 0$ . In particular, we have*

$$\mathcal{SP}(n, r) = \{\xi \in \mathcal{SP}(r) \mid l(\xi) \leq n\}.$$

*Proof.* By (6.2.1) and (6.2.3), we have  $V(n|n)_{\mathbb{K}}^{\otimes r}$  is isomorphic to a direct sum of  $\mathcal{H}_{r,\mathbb{K}}^c y_\lambda$  with  $\lambda \in \Lambda(n, r)$ . By [3, Lem. 3.50], there exist some nonnegative integers  $k_{\xi, \lambda}$  with  $k_{\xi, \lambda} \neq 0$  such that<sup>4</sup>

$$\mathcal{H}_{r,\mathbb{K}}^c y_\lambda \cong \bigoplus_{\xi \in \mathcal{SP}(r), \xi \supseteq \lambda} k_{\xi, \lambda} U^\xi, \quad (7.5.1)$$

where  $\supseteq$  is the dominance order. Since  $\xi \supseteq \lambda$  implies  $l(\xi) \leq l(\lambda) = n$ , it follows that  $V(\xi) = 0$  if  $l(\xi) > n$ , proving the first assertion. To prove the last assertion, we write  $U^\xi = \mathcal{H}_{r,\mathbb{K}}^c \epsilon$  for some idempotent  $\epsilon$ . Then, as a left  $\mathcal{Q}_v(n, r)_{\mathbb{K}}$ -module,

$$V(\xi) \cong \epsilon V(n|n)_{\mathbb{K}}^{\otimes r} \cong (V(n|n)_{\mathbb{K}}^{\otimes r})^\tau \epsilon \cong \mathcal{T}^c(n, r)_{\mathbb{K}} \otimes_{\mathcal{H}_{r,\mathbb{K}}^c} U^\xi.$$

The last assertion follows from the fact  $k_{\xi, \lambda} \neq 0$ . □

We now use Green's [14, Theorem (6.2g)] to get the following classification theorem.

<sup>3</sup>An irreducible supermodule  $X$  over an  $\mathbb{F}$ -superalgebra  $\mathcal{A}$  is called split irreducible if  $X \otimes_{\mathbb{F}} \mathbb{E}$  is irreducible over  $\mathcal{A}_{\mathbb{E}} = \mathcal{A} \otimes_{\mathbb{F}} \mathbb{E}$  for any field extension  $\mathbb{E} \supseteq \mathbb{F}$ .

<sup>4</sup>Though the result there is for the specialisation  $(\mathcal{H}_r^c)_{\mathbb{C}}$  of  $\mathcal{H}_r^c$  at  $q = 1$ . However, by the category equivalence between the module categories, the proof there carries over.

**Theorem 7.6.** *For positive integers  $n, r$ , the set*

$$\{\mathcal{T}^c(n, r)_{\mathbb{K}} \otimes_{\mathcal{H}_{r, \mathbb{K}}^c} U^\xi \cong V(\xi) \mid \xi \in \mathcal{SP}(n, r)\}$$

*forms a complete set of nonisomorphic irreducible  $\mathcal{Q}_v(n, r)_{\mathbb{K}}$ -supermodules.*

*Proof.* The case for  $n \geq r$  is seen above. We now assume  $n < r$ . Consider the natural embedding

$$\Lambda(n, r) \hookrightarrow \Lambda(r, r), \quad \lambda \mapsto \bar{\lambda} = (\lambda_1, \dots, \lambda_n, 0, \dots, 0).$$

For  $\lambda \in \Lambda(n, r)$ , let  $D(\lambda) = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_n, 0, \dots, 0) \in M_r(\mathbb{N})$  and set

$$e = \sum_{\lambda \in \Lambda(n, r)} \phi_{(D(\lambda)|0)}.$$

Then we have

$$\mathcal{Q}_v(n, r)_{\mathbb{K}} \cong \text{End}_{\mathcal{H}_{r, \mathbb{K}}^c} \left( \bigoplus_{\lambda \in \Lambda(n, r)} x_\lambda \mathcal{H}_{r, \mathbb{K}}^c \right) = e \text{End}_{\mathcal{H}_{r, \mathbb{K}}^c} \left( \bigoplus_{\mu \in \Lambda(r, r)} x_\mu \mathcal{H}_{r, \mathbb{K}}^c \right) e = e \mathcal{Q}_v(r, r)_{\mathbb{K}} e. \quad (7.6.1)$$

Meanwhile, we have

$$\begin{aligned} e(\mathcal{T}^c(r, r)_{\mathbb{K}} \otimes_{\mathcal{H}_{r, \mathbb{K}}^c} U^\xi) &= e \mathcal{T}^c(r, r)_{\mathbb{K}} \otimes_{\mathcal{H}_{r, \mathbb{K}}^c} U^\xi = e \left( \bigoplus_{\mu \in \Lambda(r, r)} x_\mu \mathcal{H}_{r, \mathbb{K}}^c \right) \otimes_{\mathcal{H}_{r, \mathbb{K}}^c} U^\xi \\ &= \left( \bigoplus_{\lambda \in \Lambda(n, r)} x_\lambda \mathcal{H}_{r, \mathbb{K}}^c \right) \otimes_{\mathcal{H}_{r, \mathbb{K}}^c} U^\xi = \mathcal{T}^c(n, r)_{\mathbb{K}} \otimes_{\mathcal{H}_{r, \mathbb{K}}^c} U^\xi. \end{aligned}$$

Since  $\{\mathcal{T}^c(r, r)_{\mathbb{K}} \otimes_{\mathcal{H}_{r, \mathbb{K}}^c} U^\xi \mid \xi \in \mathcal{SP}_r\}$  is a complete set of nonisomorphic simple  $\mathcal{Q}_v(r, r)_{\mathbb{K}}$ -supermodules, (7.6.1) together with [14, (6.2g)] implies that the set

$$\{\mathcal{T}^c(n, r)_{\mathbb{K}} \otimes_{\mathcal{H}_{r, \mathbb{K}}^c} U^\xi \mid \xi \in \mathcal{SP}_r\} \setminus \{0\} = \{\mathcal{T}^c(n, r)_{\mathbb{K}} \otimes_{\mathcal{H}_{r, \mathbb{K}}^c} U^\xi \mid \xi \in \mathcal{SP}(n, r)\}$$

is a complete set of nonisomorphic simple  $\mathcal{Q}_v(n, r)_{\mathbb{K}}$ -supermodules.  $\square$

We remark that, for the  $q$ -Schur superalgebras  $\mathcal{S}_v(m|n, r)_{\mathbb{F}}$  of type  $\mathbf{M}$  with  $m + n \geq r$ , their irreducible representations at (odd) roots of unity have been classified in [10]. It would be interesting to classify irreducible  $\mathcal{Q}_v(n, r)_{\mathbb{F}}$ -modules at roots of unity.

## REFERENCES

- [1] J. Brundan and A. Kleshchev, *Hecke-Clifford superalgebras, crystals of type  $A_{2l}^{(2)}$  and modular branching rules for  $\widehat{S}_n$* , Represent. Theory **5** (2001), 317–403.
- [2] R. Carter, *Finite groups of Lie type: conjugacy classes and complex characters*, John Wiley & Sons, New York, 1989.
- [3] S.-J. Cheng and W. Wang, *Dualities and Representations of Lie Superalgebras*, Graduate Studies in Mathematics **144**, American Mathematical Society, Providence, RI, 2012.
- [4] B. Deng, J. Du and Q. Fu, *A double Hall algebra approach to affine quantum Schur-Weyl theory*, LMS Lect. Notes Ser. **401**, 2012, Cambridge University Press, Cambridge.
- [5] B. Deng, J. Du, B. Parshall and J. Wang, *Finite dimensional algebras and quantum groups*, Mathematical Surveys and Monographs Volume **150**, Amer. Math. Soc., Providence 2008.
- [6] R. Dipper and G. James, *Representations of Hecke algebras of general linear groups*, Proc. London Math. Soc. **52** (1986), 20–52.
- [7] R. Dipper and G. James, *The  $q$ -Schur algebra*, Proc. London Math. Soc. **3** (1989), 23–50.
- [8] J. Du, *The modular representation theory of  $q$ -Schur algebras*, Trans. Amer. Math. Soc., **329** (1992), 253–271.

- [9] J. Du and H. Gu, *A realisation of the quantum supergroup  $U(\mathfrak{gl}_{m|n})$* , J. Algebra, **404** (2014), 60–99.
- [10] J. Du, H. Gu and J.-p. Wang, *Irreducible representations of  $q$ -Schur superalgebras at a root of unity*, J. Pure Appl. Algebra **218** (2014), 2012–2059.
- [11] J. Du and H. Rui, *Quantum Schur superalgebras and Kazhdan-Lusztig combinatorics*, J. Pure Appl. Algebra **215** (2011), 2715–2737.
- [12] J. Du and J. Wan, *Presenting queer Schur superalgebras*, Int. Math. Res. Not. IMRN 2015, no. 8, 2210–2272.
- [13] H. El Turkey and J. Kujawa, *Presenting Schur superalgebras*, Pacific J. Math. **262** (2013), 285–316.
- [14] J.A. Green, *Polynomial representations of  $GL_n$* , 2nd ed., with an appendix on Schensted correspondence and Littelmann paths by K. Erdmann, J. Green and M. Schocker. Lecture Notes in Mathematics **830**, Springer, Berlin, 2007.
- [15] M. Jimbo, *A  $q$ -analogue of  $U(\mathfrak{gl}(N+1))$ , Hecke algebras, and the Yang-Baxter equation*, Lett. Math. Phys. **11** (1986), 247–252.
- [16] A. Jones and M. Nazarov, *Affine Sergeev algebra and  $q$ -analogues of the Young symmetrizers for projective representations of the symmetric group*, Proc. London. Math. Soc. **78** (1999), 481–512.
- [17] T. Józefiak, *A class of projective representations of hyperoctahedral groups and Schur  $Q$ -functions*, Topics in Algebra, Banach Center Publ. **26**, Part 2, PWN-Polish Scientific Publishers, Warsaw (1990), 317–326.
- [18] A. Kleshchev, *Linear and Projective Representations of Symmetric Groups*, Cambridge Tracts in Mathematics **136**, CUP 2005.
- [19] H. Mitsuhashi, *Schur-Weyl reciprocity between the quantum superalgebra and the Iwahori-Hecke algebra*, Algebr. Represent. Theory **9** (2006), 309–322.
- [20] G.I. Olshanski, *Quantized universal enveloping superalgebra of type  $Q$  and a super-extension of the Hecke algebra*, Lett. Math. Phys. **24** (1992), 93–102.
- [21] A. Sergeev, *Tensor algebra of the identity representation as a module over the Lie superalgebras  $GL(n, m)$  and  $Q(n)$* , Math. USSR Sbornik **51** (1985), 419–427.
- [22] J. Wan and W. Wang, *Frobenius character formula and spin generic degrees for Hecke-Clifford algebra*, Proc. London Math. Soc. **106** (2013), 287–317.
- [23] J. Wan and W. Wang, *Spin Kostka polynomials*, J. Algebraic Combin. **37** (2013), 117–138.

(DU) SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF NEW SOUTH WALES, UNSW SYDNEY 2052, AUSTRALIA.

*E-mail address:* j.du@unsw.edu.au

(WAN) SCHOOL OF MATHEMATICS AND STATISTICS, BEIJING INSTITUTE OF TECHNOLOGY, BEIJING, 100081, P.R. CHINA.

*E-mail address:* wjk302@hotmail.com